





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 401/12, 403/12, A61K 31/55, C07D 223/16, 471/04, 413/12, 405/12, 409/12, 417/12		A1	(11) International Publication Number: WO 00/21951
			(43) International Publication Date: 20 April 2000 (20.04.00)
(21) International Application Number: PCT/EP99/07763		Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). VONG, Antonio, Kuok, Keong [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). (74) Agent: GARRETT, Michael; SmithKline Beecham, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).	
(22) International Filing Date: 6 October 1999 (06.10.99)			
(30) Priority Data:			
9821976.9 8 October 1998 (08.10.98) GB 9824340.5 6 November 1998 (06.11.98) GB 9910711.2 7 May 1999 (07.05.99) GB 9918032.5 30 July 1999 (30.07.99) GB			
(71) Applicant (for all designated States except US): SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(72) Inventors; and			
(75) Inventors/Applicants (for US only): HADLEY, Michael, Stewart [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). JOHNSON, Christopher, Norbert [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). MACDONALD, Gregor, James [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). STEMP, Geoffrey [GB/GB]; SmithKline			
		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	

(54) Title: TETRAHYDROBENZAZEPINE DERIVATIVES USEFUL AS MODULATORS OF DOPAMINE D₃ RECEPTORS (ANTI-PSYCHOTIC AGENTS)



(57) Abstract

Compounds of formula (I), wherein R² represents a hydrogen atom or a C₁₋₄alkyl group; q is 1 or 2; A represents a group of formula (a), (b), (c) or (d), wherein Ar represents an optionally substituted phenyl ring or an optionally substituted 5- or 6-membered aromatic heterocyclic ring; or an optionally substituted bicyclic ring system; Ar¹ and Ar² each independently represent an optionally substituted phenyl ring or an optionally substituted 5- or 6-membered aromatic heterocyclic ring; and Y represents a bond, —NHCO—, —CONH—, —CH₂—, or —(CH₂)_mY¹(CH₂)_n—, wherein Y¹ represents O, S, SO₂, or CO and m and n each represent zero or 1 such that the sum of m+n is zero or 1; providing that when A represents a group of formula (a), any substituent present in Ar *ortho* to the carboxamide moiety is necessarily a hydrogen or a methoxy group; r and s independently represent an integer from zero to 3 such that the sum of r and s is equal to an integer from 1 to 4; V represents a bond, O or S; and salts thereof. Compounds of formula (I) and their salts have affinity for dopamine receptors, in particular the D₃ receptor, and thus potential in the treatment of conditions wherein modulation of the D₃ receptor is beneficial, e.g. as antipsychotic agents.

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakistan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						


TETRAHYDROBENZAZEPINE DERIVATIVES USEFUL AS MODULATORS OF DOPAMINE D₃ RECEPTORS (ANTI-PSYCHOTIC AGENTS)

The present invention relates to novel tetrahydrobenzazepine derivatives, processes for their preparation, pharmaceutical compositions containing them and their use in therapy, as modulators of dopamine D₃ receptors, in particular as antipsychotic agents.

US Patent No. 5,294,621 describes tetrahydropyridine derivatives of the formula:




10





wherein is an optionally substituted thienyl or optionally substituted phenyl ring; R¹, R² and R³ are each *inter alia* hydrogen; X is *inter alia* (CH₂)_mNR⁷CO; m is 2-4; and Ar¹ is an optionally substituted heterocyclic ring or an optionally substituted phenyl ring. The compounds are said to be useful as antiarrhythmic agents.

15


EPA 431,580 describes compounds of formula



wherein R is OR³, NR⁴R⁵, or N(OR⁴)R⁵, R⁴ and R⁵ are *inter alia* hydrogen, lower alkyl, aroyl or heteroaroyl; m is zero, 1 or 2; R¹ is hydrogen, aryl or various heteroaryl groups; n is zero or 1-4; and R² is:



or



25


The compounds are said to be dopaminergic agents useful as antipsychotics, antihypertensives and also of use in the treatment of hyperprolactinaemia-related conditions and several central nervous system disorders.

WO 95/10513 describes benzothiophene derivatives and related compounds as estrogen agonists.

WO 97/43262 and WO 98/06699 describe tetrahydroisoquinoline derivatives as having affinity for the dopamine D₃ receptor.

5 We have now found a class of tetrahydrobenzazepine derivatives which have affinity for dopamine receptors, in particular the D₃ receptor, and thus potential in the treatment of conditions wherein modulation of the D₃ receptor is beneficial, eg as antipsychotic agents.

In a first aspect the present invention provides compounds of formula (I) :



Formula (I)

wherein:

R¹ represents a substituent selected from: a hydrogen or halogen atom; a hydroxy, cyano, nitro, trifluoromethyl, trifluoromethoxy, trifluoromethanesulfonyloxy, pentafluoroethyl, C₁₋₄alkyl, C₁₋₄alkoxy, arylC₁₋₄alkoxy, C₁₋₄alkylthio, C₁₋₄alkoxyC₁₋₄alkyl, C₃₋₆cycloalkylC₁₋₄alkoxy, C₁₋₄alkanoyl, C₁₋₄alkoxycarbonyl, C₁₋₄alkylsulfonyl, C₁₋₄alkylsulfonyloxy, C₁₋₄alkylsulfonylC₁₋₄alkyl, arylsulfonyl, arylsulfonyloxy, arylsulfonylC₁₋₄alkyl, C₁₋₄alkylsulfonamido, C₁₋₄alkylamido, C₁₋₄alkylsulfonamidoC₁₋₄alkyl, C₁₋₄alkylamidoC₁₋₄alkyl, arylsulfonamido, arylcarboxamido, arylsulfonamidoC₁₋₄alkyl, arylcarboxamidoC₁₋₄alkyl, aroyl, aroylC₁₋₄alkyl, or arylC₁₋₄alkanoyl group; a group R³OCO(CH₂)_p, R³CON(R⁴)(CH₂)_p, R³R⁴NCO(CH₂)_p or R³R⁴NSO₂(CH₂)_p where each of R³ and R⁴ independently represents a hydrogen atom or a C₁₋₄alkyl group or R³R⁴ forms part of a C₃₋₆azacycloalkane or C₃₋₆(2-oxo)azacycloalkane ring and p represents zero or an integer from 1 to 4; or a group Ar³-Z, wherein Ar³ represents an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring and Z represents a bond, O, S, or CH₂;


R² represents a hydrogen atom or a C₁₋₄alkyl group;

q is 1 or 2;


30 A represents a group of the formula (a), (b) (c) or (d):




(a)



(b)



(c)



(d)

wherein

35 Ar represents an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring; or an optionally substituted bicyclic ring system;

Ar¹ and Ar² each independently represent an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring; and

Y represents a bond, -NHCO-, -CONH-, -CH₂-, or -(CH₂)_mY¹(CH₂)_n-, wherein Y¹ represents O, S, SO₂, or CO and m and n each represent zero or 1 such that the sum of m+n is zero or 1; providing that when A represents a group of formula (a), any substituent present in Ar *ortho* to the carboxamide moiety is necessarily a hydrogen or a methoxy group;

r and s independently represent an integer from zero to 3 such that the sum of r and s is equal to an integer from 1 to 4;

V represents a bond, O or S;
and salts thereof.

In the compounds of formula (I) above an alkyl group or moiety may be straight or branched. Alkyl groups which may be employed include methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl and any branched isomers thereof such as isopropyl, t-butyl, sec-butyl, and the like.

When R¹ represents an arylC₁₋₄alkoxy, arylsulfonyl, arylsulfonyloxy, arylsulfonylC₁₋₄alkyl, arylsulfonamido, arylcarboxamido, arylsulfonamidoC₁₋₄alkyl, arylcarboxamidoC₁₋₄alkyl, aroyl, aroylC₁₋₄alkyl, or arylC₁₋₄alkanoyl group, the aryl moiety may be selected from an optionally substituted phenyl ring or an optionally substituted 5- or 6-membered heterocyclic ring. In the group R¹ an aryl moiety may be optionally substituted by one or more substituents selected from hydrogen, halogen, amino, cyano, C₁₋₄alkyl, C₁₋₄alkylamino, C₁₋₄dialkylamino, C₁₋₄alkylamido, C₁₋₄alkanoyl, or R⁵R⁶NCO where each of R⁵ and R⁶ independently represents a hydrogen atom or C₁₋₄alkyl group.

A halogen atom present in the compounds of formula (I) may be fluorine, chlorine, bromine or iodine.

When q is 2, the substituents R¹ may be the same or different.

An optionally substituted 5- or 6-membered heterocyclic aromatic ring, as defined for any of the groups Ar, Ar¹, Ar² or Ar³ may contain from 1 to 4 heteroatoms selected from O, N or S. When the ring contains 2-4 heteroatoms, one is preferably selected from O, N and S and the remaining heteroatoms are preferably N. Examples of 5 and 6-membered heterocyclic groups include furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, pyridyl, triazolyl, triazinyl, pyridazyl, pyrimidinyl, pyrazolyl, isothiazolyl, and isoxazolyl.

Examples of bicyclic, for example bicyclic aromatic or heteroaromatic, ring systems for Ar include naphthyl, indazolyl, indolyl, benzofuranyl, benzothienyl, benzothiazolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzisothiazolyl, quinolinyl, quinoxolinyl, quinazolinyl, cinnolinyl, isoquinolinyl, pyrazolo[1,5-a]pyrimidyl, pyrrolo[3,2-b]pyridyl, pyrrolo[3,2-c]pyridyl, thieno[3,2-b]thiophenyl, 1,2-dihydro-2-oxo-quinolinyl, 3,4-dihydro-3-oxo-2H-benzoxazinyl, 1,2-dihydro-2-oxo-3H-indolyl.

The rings Ar, Ar¹, or Ar² may each independently be optionally substituted by one or more substituents selected from: a hydrogen or halogen atom, or a hydroxy, oxo, cyano, nitro, trifluoromethyl, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkylenedioxy, C₁₋₄alkanoyl,

C₁₋₄alkylsulfonyl, C₁₋₄alkylsulfinyl, C₁₋₄alkylthio, R⁷SO₂N(R⁸)-, R⁷R⁸NSO₂-, R⁷R⁸N-, R⁷R⁸NCO-, or R⁷CON(R⁸)- group wherein each of R⁷ and R⁸ independently represents a hydrogen atom or a C₁₋₄ alkyl group, or R⁷R⁸ together form a C₃₋₆ alkylene chain.

5 Alternatively, Ar and Ar² may be optionally substituted by one or more 5- or 6-membered heterocyclic rings, as defined above, optionally substituted by a C₁₋₂ alkyl or R⁷R⁸N- group; wherein R⁷ and R⁸ are as defined above.

In the rings Ar and Ar² substituents positioned *ortho* to one another may be linked to form a 5- or 6- membered ring.

10 It will be appreciated that for use in medicine the salts of formula (I) should be physiologically acceptable. Suitable physiologically acceptable salts will be apparent to those skilled in the art and include for example acid addition salts formed with inorganic acids eg. hydrochloric, hydrobromic, sulfuric, nitric or phosphoric acid; and organic acids eg. succinic, maleic, acetic, fumaric, citric, tartaric, benzoic, p-toluenesulfonic, 15 methanesulfonic or naphthalenesulfonic acid. Other non-physiologically acceptable salts eg. oxalates, may be used, for example in the isolation of compounds of formula (I) and are included within the scope of this invention. Also included within the scope of the invention are solvates and hydrates of compounds of formula (I).

20 Certain of the compounds of formula (I) may form acid addition salts with one or more equivalents of the acid. The present invention includes within its scope all possible stoichiometric and non-stoichiometric forms.

The compounds of formula (I) can exist in the form of *cis*- and *trans*- isomers with respect to the configuration at the cyclohexyl ring. When A represents a group (c) the compounds may also exist as geometric isomers around the double bond. The present 25 invention includes within its scope all such isomers, including mixtures. Preferably the compounds of the invention are in the *trans* configuration with respect to the cyclohexyl ring. For compounds of formula (I) where A represents a group (c), *trans* geometry of the double bond is preferred.

30 In compounds of formula (I), it is preferred that R¹ represents a substituent selected from: a halogen atom, methyl, cyano, acetyl, trifluoromethyl, pentafluoroethyl, methylsulphonyl, methylsulphonyloxy or trifluoromethoxy group. Alternatively, it is preferred that R¹ represents a group Ar³Z, where Z is a bond and Ar³ is a 5- or 6-membered ring heterocycle, optionally substituted by a methyl group, containing at least one N and one O atom. Preferably q is 1. R² is preferably a hydrogen atom.

35 When the group A is a group of formula (a), preferred examples of Ar include optionally substituted phenyl, indolyl, pyrazolo[1,5-a]pyrimidyl, cinnolinyl, quinolinyl, benzo[b]furanyl or pyrrolopyridyl.

40 When the group A is a group of formula (b), preferred examples of Ar¹ include optionally substituted phenyl, Y is preferably a bond, and preferred examples of Ar² include optionally substituted phenyl, pyridyl, pyrimidinyl, isoxazolyl, oxazolyl or oxadiazolyl.

When the group A is a group of formula (c), preferred examples of Ar include optionally substituted phenyl.

It is also preferred that the rings Ar, Ar¹, or Ar² are each independently optionally substituted by one or more substituents selected from: a hydrogen or halogen atom, cyano, methoxy, trifluoromethyl, methylenedioxy, acetyl, acetylamino, methylsulfonyl, methylsulfonyloxy, methylaminosulfonyl, methylsulfonylamino, or methylaminocarbonyl group.

Certain of the substituted heteroaromatic ring systems included in compounds of formula (I) may exist in one or more tautomeric forms. The present invention includes within its scope all such tautomeric forms, including mixtures.

Particular compounds according to the invention include those specifically exemplified and named hereinafter:-

trans-3-(2-(1-(4-(4-Quinoliny)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-(*E*)-3-(2-(1-(4-(3-(3-Methylsulfonyl)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-(*E*)-3-(2-(1-(4-(3-(4-Fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-3-(2-(1-(4-(2-Indolyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-3-(2-(1-(4-(3-(3-Pyridyl)phenyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-3-(2-(1-(4-Phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-3-(2-(1-(4-(3-Indolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-3-(2-(1-(4-(4-Quinoliny)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-(*E*)-3-(2-(1-(4-(3-(4-Fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-6-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-6-Methoxy-3-(2-(1-(4-(4-quinoliny)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-6-Methoxy-3-(2-(1-(4-(3-pyrrolo[2,3-*b*]pyridyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-(*E*)-7-Cyano-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-7-Cyano-3-(2-(1-(4-(3-pyrrolo[2,3-*b*]pyridyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-7-Cyano-3-(2-(1-(4-(3-(3-(5-methyl)-1,2,4-oxadiazolyl)benzoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-(*E*)-7-Cyano-3-(2-(1-(4-(5-quinoliny)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

trans-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-acetylamino)phenylpropenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

- trans*-7-Cyano-3-(2-(1-(4-(6-(3,4-dihydro-3-oxo)-2*H*-benzoxazinyl)carboxamido)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(6-(1,2-dihydro-2-oxo)quinolinyl)propenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 5 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-fluoro-4-acetyl amino)phenylpropenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(8-(1,2-dihydro-2-oxo)quinolinyl)propenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(5-(8-fluoro)quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 10 *trans*-7-Acetyl-3-(2-(1-(4-(5-quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-3-(2-(1-(4-(5-(2-Methyl)quinolinyl)carboxamido)cyclohexyl)ethyl)-7-methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 15 *trans*-3-(2-(1-(4-(3-(3-(5-Methyl)-1,2,4-oxadiazolyl)benzoyl)amino)cyclohexyl)ethyl)-7-methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-pyrrolo[2,3-*b*]pyridyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 20 *trans*-(*E*)-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-(5-(3-Methyl)isoxazolyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 25 *trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(4-fluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(2,5-difluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(2-naphthylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 30 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2,4-difluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2,5-difluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 35 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(3-phenylpropanoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 40 *trans*-7-Cyano-3-(2-(1-(4-(8-(1,4-dihydro-4-oxo)quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(2-naphthyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-methoxy)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-methoxy)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 5 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(4-methoxy)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-acetyl)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(4-acetyl)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 10 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-cyano)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-cyano)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 15 *trans*-7-Cyano-3-(2-(1-(4-(3-(5-(3-methyl)isoxazolyl)benzoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(7-(1,2-dihydro-2-oxo)quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*Z*)-7-Cyano-3-(2-(1-(4-(3-phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 20 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-pyridyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(1-(4-fluoro)naphthyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 25 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(6-benzodioxanyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-(5-fluoro)indolyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-(1-methyl)benzimidazolyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 30 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(7-benzofuranyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(5-(3-methyl)indolyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 35 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(6-(2,3-dihydro-2-oxo)indolyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(2-benzofuranylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(4-(2-methyl)indolyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 40 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(5-benzimidazolyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2,3-methylenedioxy)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-(1-(2-oxo)pyrrolidinyl))phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 5 *trans*-7-Cyano-3-(2-(1-(4-(2-indolylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(2-benzothiophenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-(3-bromo)thiophenyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 10 *trans*-7-Cyano-3-(2-(1-(4-(3-(2-pyridyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(3-(5-pyrimidinyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 15 *trans*-7-Cyano-3-(2-(1-(4-(3-(4-cyanophenyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(3-(3-(5-ethyl)-1,2,4-oxadiazolyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-thiophenyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 20 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-furanyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-thiophenyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 25 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-furanyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(4-quinolinyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(5-pyrimidinyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 30 *trans*-7-Cyano-3-(2-(1-(4-(2,4-difluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(1-naphthyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 35 *trans*-7-Acetyl-3-(2-(1-(4-(3-pyrrolo[2,3-*b*]pyridyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Acetyl-3-(2-(1-(4-(4-fluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Acetyl-3-(2-(1-(4-(3-(3-(5-methyl)-1,2,4-oxadiazolyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 40 *trans*-(*E*)-7-Acetyl-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(6-(2-amino)benzothiazolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

- trans*-7-Cyano-3-(2-(1-(4-(6-(2-methyl)benzothiazolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(6-(2,3-dihydro-2-oxo)indolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 5 *trans*-7-Cyano-3-(2-(1-(4-(5-(2,3-dihydro-2-oxo)indolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(4-methylaminocarbonyl)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(5-(2-amino)benzoxazolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 10 *trans*-7-Cyano-3-(2-(1-(4-(6-(1,2-dihydro-2-oxo)quinolinyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(7-(1,2-dihydro-2-oxo)quinolinyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 15 *trans*-(*E*)-7-Acetyl-3-(2-(1-(4-(3-(3-methoxy)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Acetyl-3-(2-(1-(4-(3-(2-cyano)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Acetyl-3-(2-(1-(4-(3-(3-thiophenyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 20 *trans*-(*E*)-7-Acetyl-3-(2-(1-(4-(3-(8-(1,2-dihydro-2-oxo)quinolinyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(3-(1-pyrazolyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 25 *trans*-7-Cyano-3-(2-(1-(4-(2-thiophenyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(3-benzothiophenyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(3-(2-(5-methyl)-1,3,4-oxadiazolyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 30 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-naphthyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Acetyl-3-(2-(1-(4-(3-benzothiophenyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 35 *trans*-(*E*)-7-Acetyl-3-(2-(1-(4-(3-(4-acetamido)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Acetyl-3-(2-(1-(4-(6-(2-amino)benzothiazolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Acetyl-3-(2-(1-(4-(8-(1,4-dihydro-4-oxo)quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 40 *trans*-(*E*)-7-Acetyl-3-(2-(1-(4-(3-(2-acetyl)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Acetyl-3-(2-(1-(4-(2-benzothiophenyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(5-(3-acetyl)indolyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(3-(5-(3-methyl)-1,2,4-oxadiazolyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 5 *trans*-7-Cyano-3-(2-(1-(4-(5-(2-methyl)benzimidazolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(6-quinoxaliny)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-(2-acetyl)furanyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 10 *trans*-7-Cyano-3-(2-(1-(4-(6-(2-amino)benzoxazolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(6-(3,4-dihydro-2-oxo)-2*H*-benzoxazinyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 15 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-fluoro-5-acetamido)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(2-benzothiophenyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-thienyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 20 *trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(5-quinoliny)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-(5-Methyl)-1,2,4-oxadiazolyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 25 *trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(8-(1,4-dihydro-4-oxo)quinoliny)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-acetamido-2-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 30 *trans*-(*E*)-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-acetyl)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-fluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 35 *trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(2,4-difluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(2-naphthyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(7-(3,4-dihydro-3-oxo)-2*H*-benzoxazinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 40 *trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(5-(2-methyl)quinoliny)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(2-fluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;


- trans*-3-(2-(1-(4-(5-(2-Methyl)quinolinyl)carboxamide)cyclohexyl)ethyl)-7-methanesulphonyloxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(2-indolyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 5 *trans*-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(2-benzothiophenyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-thienyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(5-quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 10 *trans*-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-pyrrolo[2,3-*b*]pyridyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(8-(1,4-dihydro-4-oxo)quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 15 *trans*-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-(5-Methyl)-1,2,4-oxadiazolyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(2-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 20 *trans*-3-(2-(1-(4-(5-(2-Methyl)quinolinyl)carboxamide)cyclohexyl)ethyl)-7-methanesulphonyloxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-7-methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 25 *trans*-3-(2-(1-(4-(3-(2-(4-Methyl)oxazolyl)benzoyl)amino)cyclohexyl)ethyl)-7-methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-3-(2-(1-(4-(3-(trifluoromethyl)benzoyl)amino)cyclohexyl)ethyl)-7-methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-3-(2-(1-(4-(5-(8-Chloro-2-methyl)quinolinyl)carboxamide)cyclohexyl)ethyl)-7-methanesulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 30 *trans*-7-(5-(3-Methyl)isoxazolyl)-3-(2-(1-(4-(2-fluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-(3-(5-Methyl)isoxazolyl)-3-(2-(1-(4-(4-fluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 35 *trans*-(*E*)-7-(2-(5-Methyl)oxazolyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-(3-(5-Methyl)isoxazolyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-(2-Pyridyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 40 *trans*-(*E*)-7-(2-Pyrimidyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-(1-Pyrrolidinylcarbonyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 45 *trans*-7-(1-Pyrrolidinylcarbonyl)-3-(2-(1-(4-(3-pyrrolo[2,3-*b*]pyridyl)

- carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-(5-Pyrimidyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)
 cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(5-Pyrimidinyl)-3-(2-(1-(4-(3-(3-(5-Methyl)-1,2,4-
 5 oxadiazolyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(5-Pyrimidinyl)-3-(2-(1-(4-(5-(2-methyl)quinolinyl)carboxamido)cyclohexyl)
 ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(3-(5-Methyl)isoxazolyl)-3-(2-(1-(4-(5-(2-methyl)quinolinyl)carboxamido)
 cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-(2-Pyridyl)-3-(2-(1-(4-(3-(2-cyano)phenylpropenoyl)amino)
 cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-(2-Pyridyl)-3-(2-(1-(4-(3-(3-cyano)phenylpropenoyl)amino)
 cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-(2-Pyridyl)-3-(2-(1-(4-(3-(4-cyano)phenylpropenoyl)amino)
 cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-3-(2-(1-(4-(5-(8-Fluoro-2-methyl)quinolinyl)carboxamido)cyclohexyl)ethyl)-7-
 methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-3-(2-(1-(4-(5-(8-Fluoro-2-methyl)quinolinyl)carboxamido)cyclohexyl)ethyl)-7-
 methylsulphonyloxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-3-(2-(1-(4-(3-(2-(5-Methyl)oxazolyl)benzoyl)amino)cyclohexyl)ethyl)-7-
 methylsulphonyloxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

These compounds may be in the form of their free base or physiologically acceptable salts thereof, particularly the monohydrochloride or monomesylate salts.

The present invention also provides a process for preparing compounds of formula (I) which process comprises :

(a) reacting a compound of formula (II):



Formula (II)

wherein R¹, R² and q are as hereinbefore defined, with a compound of formula (III):




Formula (III)

wherein A is as hereinbefore defined and X is a halogen atom or the residue of an activated ester;

(b) to prepare a compound of formula (I) by reacting a compound of formula (II) with a compound A-Br, or A-I, or A-OSO₂CF₃ in the presence of carbon monoxide and a catalyst such as *trans*-bis-triphenylphosphinepalladium(II)bromide;

(c) to prepare a compound of formula (I) wherein R¹ is Ar³-Z and Z is a bond,
5 reacting a compound of formula (IV):




Formula (IV)

10

wherein R² and A are as hereinbefore defined and one R^{1a} represents a group W wherein W is a halogen atom or a trifluoromethylsulfonyloxy group, or W is a group M selected from a boron derivative e.g. a boronic acid function B(OH)₂ or a metal function such as trialkylstannyl e.g. SnBu₃, zinc halide or magnesium halide, and when q is 2 the other R^{1a} is R¹; with a compound Ar³-W¹, wherein W¹ is a halogen atom or a trifluoromethylsulfonyloxy group when W is a group M or W¹ is a group M when W is a halogen atom or a trifluoromethylsulfonyloxy group;

15

(d) to prepare a compound of formula (I) wherein R¹ is Ar³-Z and Z is O or S, reacting a compound of formula (V):




20

Formula (V)

wherein R² and A are as hereinbefore defined and one R^{1b} represents a group ZH and when q is 2 the other R^{1b} represents R¹; with a reagent serving to introduce the group Ar³;

25

(e) to prepare a compound of formula (I) where Y is a bond, reaction of a compound of formula (VI):



30

Formula (VI)

wherein R^1 , R^2 , Ar^1 , W and q are as hereinbefore defined, with a compound Ar^2-W^1 , wherein W^1 is a halogen atom or a trifluoromethylsulfonyloxy group when W is a group M , or W^1 is a group M when W is a halogen atom or a trifluoromethylsulfonyloxy group.

5 (f) interconversion of one compound of formula (I) to a different compound of formula (I) e.g. (i) alkylation of a compound (I) wherein R^2 represents hydrogen, (ii) conversion of one R^1 from alkoxy (e.g. methoxy) to hydroxy, or (iii) conversion of R^1 from hydroxy to sulfonyloxy, eg alkylsulfonyloxy or trifluoromethanesulfonyloxy; (iv) conversion of a compound wherein Y represents S to a compound wherein Y is SO_2 or
10 (v) conversion of Y from CO to CH_2 ;

(g) separation of *cis*- and *trans*- isomers of compounds of formula (I) by conventional methods, e.g. chromatography or crystallisation; and optionally thereafter forming a salt of formula (I).


Process (a) may be effected using conventional methods for the formation of an amide bond. When X is the residue of an activated ester this may be formed with e.g. a carbodiimide such as 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide. The reaction may be carried out in a solvent such as dichloromethane.

Reaction of a compound of formula (IV) with Ar^3W^1 , according to process (c) or a compound of formula (VI) with Ar^2-W^1 according to process (e) may be effected in the
20 presence of a transition metal eg palladium catalyst such as *bis*-triphenylphosphinepalladium dichloride or *tetrakis*-triphenylphosphinepalladium (0). When M represents a boronic acid function such as $B(OH)_2$ the reaction may be carried out under basic conditions, for example using aqueous sodium carbonate in a suitable solvent such as dioxane. When M is trialkylstannyl the reaction may be carried out in an
25 inert solvent, such as xylene or dioxane optionally in the presence of $LiCl$. When M is a zinc or magnesium halide the reaction may be effected in an aprotic solvent such as tetrahydrofuran. The substituent W is preferably a halogen atom such as bromine, or a sulfonyloxy group such as trifluoromethylsulfonyloxy; and W^1 is preferably a group M , such as trialkylstannyl or $B(OH)_2$.

30 In process (d) the reagent serving to introduce the group Ar^3 is preferably a compound of formula Ar^3-Hal , wherein Hal is a halogen atom. The reaction may be effected in the presence of a base, such as potassium carbonate, in a solvent such as dimethylformamide.

35 Interconversion reactions according to process (f) may be effected using methods well known in the art.


Compounds of formula (II) may be prepared by conversion of a compound of formula (VII), wherein R^1 and q are as hereinbefore defined,



40 **Formula (VII)**


into a corresponding ketone, followed by reductive amination. This may be effected by methods well known in the art for (i) conversion of a ketal to a ketone in the presence of aqueous acid; followed by (ii) reductive amination of the ketone with R^2NH_2 or ammonium acetate in the presence of a reducing agent. Suitable reducing agents which may be employed include sodium borohydride, cyanoborohydride or triacetoxyborohydride under acidic conditions, or catalytic hydrogenation. The reaction may conveniently be effected in a solvent such as methanol, ethanol or dichloroethane.

A compound of formula (VII) may itself be prepared by reacting a compound of formula (VIII):



Formula (VIII)


wherein R^1 and q are as hereinbefore defined;
with a compound of formula (IX):



Formula (IX)

in the presence of a reducing agent. Suitable reducing agents which may be employed include sodium borohydride, cyanoborohydride or triacetoxyborohydride under acidic conditions, or catalytic hydrogenation. The reaction may conveniently be effected in a solvent such as ethanol or dichloroethane.

The individual *cis*- and *trans*- isomers of a compound of formula (II) may be prepared starting from *cis*- or *trans*- 4-amino-cyclohexaneacetic acid (T.P. Johnson, *et al.*, J. Med. Chem., 1997, (20), 279-290) followed by functional group interchange and/or protection using methods well known in the art, to give the individual *cis*- or *trans*- isomers of a compound of formula (X):




Formula (X)

wherein R^2 is as hereinbefore defined, and P is a protecting group, for example trifluoroacetyl or *tert*-butoxycarbonyl. Subsequent reaction of a compound of formula (X) with a compound of formula (VIII) in the presence of a reducing agent as described above followed by deprotection using standard methodology gives the individual isomers of a compound of formula (II) wherein R^2 is as hereinbefore defined.

Compounds of formula (III) are known or may be prepared using standard procedures.

Compounds of formula (IV), (V) or (VI) may be prepared by processes analogous to (a), (b), (c) and (d) described above. Compounds Ar^2W^1 , Ar^3W^1 and Ar^3Hal are commercially available or may be prepared by standard methods. Compounds of formula (VIII), where for example R^1 is a halogen, methoxy, acetyl, cyano, carboxylic acid or carboxamide group are known in the literature or may be prepared by known methods. The compound of formula (IX) is likewise known in the literature.

Conversion of a compound of formula (VIII) where R^1 is a cyano or acetyl group to a compound of formula (VIII) where R^1 is a group Ar^3Z , where Ar is an oxadiazole or an isoxazole ring and Z is a bond, may be carried out by (i) conversion to a compound of formula (XI), where R^1 and q are as hereinbefore defined, using standard methods; (ii) conversion of R^1 from cyano to oxadiazolyl using known methods, or conversion of acetyl to isoxazolyl using known methods; (iii) deprotection of a compound of formula (XI) to a compound of formula (VIII) using standard methods.



Formula (XI)

Compounds of formula (I) have been found to exhibit affinity for dopamine receptors, in particular the D_3 receptor, and are expected to be useful in the treatment of disease states which require modulation of such receptors, such as psychotic conditions. Compounds of formula (I) have also been found to have greater affinity for dopamine D_3 than for D_2 receptors. The therapeutic effect of currently available antipsychotic agents (neuroleptics) is generally believed to be exerted via blockade of D_2 receptors; however this mechanism is also thought to be responsible for undesirable extrapyramidal side effects (eps) associated with many neuroleptic agents. Without wishing to be bound by theory, it has been suggested that blockade of the recently characterised dopamine D_3 receptor may give rise to beneficial antipsychotic activity without significant eps. (see for example Sokoloff et al, Nature, 1990; 347: 146-151; and Schwartz et al, Clinical Neuropharmacology, Vol 16, No. 4, 295-314, 1993). Preferred compounds of the present invention are therefore those which have higher affinity for dopamine D_3 than dopamine D_2 receptors (such affinity can be measured using standard methodology for example using cloned dopamine receptors). Said compounds may advantageously be used as selective modulators of D_3 receptors.

The compounds of formula (I) are of potential use as antipsychotic agents for example in the treatment of schizophrenia, schizo-affective disorders, psychotic

depression, mania, paranoid and delusional disorders. Furthermore, they could have utility as adjunct therapy in Parkinsons Disease, particularly with compounds such as L-DOPA and possibly dopaminergic agonists, to reduce the side effects experienced with these treatments on long term use (eg see Schwartz et al., Brain Res. Reviews, 1998, 26, 236-242). From the localisation of D3 receptors, it could also be envisaged that the compounds could also have utility for the treatment of substance abuse where it has been suggested that D3 receptors are involved (eg see Levant, 1997, Pharmacol. Rev., 49, 231-252). Examples of such substance abuse include alcohol, cocaine and nicotine abuse. Other conditions which may be treated by the compounds include dyskinetic disorders such as Parkinson's disease, neuroleptic-induced parkinsonism and tardive dyskinesias; depression; anxiety, cognitive impairment including memory disorders such as Alzheimers disease, eating disorders, sexual dysfunction, sleep disorders, emesis, movement disorders, obsessive-compulsive disorders, amnesia, aggression, autism, vertigo, dementia, circadian rhythm disorders and gastric motility disorders e.g. IBS.

In a further aspect therefore the present invention provides a method of treating conditions which require modulation of dopamine D₃ receptors, for example psychoses such as schizophrenia, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I) or a physiologically acceptable salt thereof.

The invention also provides the use of a compound of formula (I) or a physiologically acceptable salt thereof in the manufacture of a medicament for the treatment of conditions which require modulation of dopamine D₃ receptors, for example psychoses such as schizophrenia.

A preferred use for D₃ antagonists according to the present invention is in the treatment of psychoses such as schizophrenia.

For use in medicine, the compounds of the present invention are usually administered as a standard pharmaceutical composition. The present invention therefore provides in a further aspect pharmaceutical compositions comprising a novel compound of formula (I) or a physiologically acceptable salt thereof and a physiologically acceptable carrier.

The compounds of formula (I) may be administered by any convenient method, for example by oral, parenteral, buccal, sublingual, nasal, rectal or transdermal administration and the pharmaceutical compositions adapted accordingly.

The compounds of formula (I) and their physiologically acceptable salts which are active when given orally can be formulated as liquids or solids, for example syrups, suspensions or emulsions, tablets, capsules and lozenges.

A liquid formulation will generally consist of a suspension or solution of the compound or physiologically acceptable salt in a suitable liquid carrier(s) for example an aqueous solvent such as water, ethanol or glycerine, or a non-aqueous solvent, such as polyethylene glycol or an oil. The formulation may also contain a suspending agent, preservative, flavouring or colouring agent.

A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and cellulose.

A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion or suspension then filled into a soft gelatin capsule.

Typical parenteral compositions consist of a solution or suspension of the compound or physiologically acceptable salt in a sterile aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration.

Compositions for nasal administration may conveniently be formulated as aerosols, drops, gels and powders. Aerosol formulations typically comprise a solution or fine suspension of the active substance in a physiologically acceptable aqueous or non-aqueous solvent and are usually presented in single or multidose quantities in sterile form in a sealed container, which can take the form of a cartridge or refill for use with an atomising device. Alternatively the sealed container may be a unitary dispensing device such as a single dose nasal inhaler or an aerosol dispenser fitted with a metering valve which is intended for disposal once the contents of the container have been exhausted. Where the dosage form comprises an aerosol dispenser, it will contain a propellant which can be a compressed gas such as compressed air or an organic propellant such as a fluorochlorohydrocarbon. The aerosol dosage forms can also take the form of a pump-atomiser.

Compositions suitable for buccal or sublingual administration include tablets, lozenges and pastilles, wherein the active ingredient is formulated with a carrier such as sugar and acacia, tragacanth, or gelatin and glycerin.

Compositions for rectal administration are conveniently in the form of suppositories containing a conventional suppository base such as cocoa butter.

Compositions suitable for transdermal administration include ointments, gels and patches.

Preferably the composition is in unit dose form such as a tablet, capsule or ampoule.

Each dosage unit for oral administration contains preferably from 1 to 250 mg (and for parenteral administration contains preferably from 0.1 to 25 mg) of a compound of the formula (I) or a physiologically acceptable salt thereof calculated as the free base.

The physiologically acceptable compounds of the invention will normally be administered in a daily dosage regimen (for an adult patient) of, for example, an oral dose of between 1 mg and 500 mg, preferably between 10 mg and 400 mg, e.g. between 10 and 250 mg or an intravenous, subcutaneous, or intramuscular dose of between 0.1 mg and 100 mg, preferably between 0.1 mg and 50 mg, e.g. between 1 and 25 mg of the compound of the formula (I) or a physiologically acceptable salt thereof calculated as the free base, the compound being administered 1 to 4 times per day. Suitably the compounds will be administered for a period of continuous therapy, for example for a week or more.

Biological Test Methods

The ability of the compounds to bind selectively to human D₃ dopamine receptors can be demonstrated by measuring their binding to cloned receptors. The inhibition constants (K_i) of test compounds for displacement of [¹²⁵I] idosulpride binding to human D₃ dopamine receptors expressed in CHO cells were determined as follows. The cell lines were shown to be free from bacterial, fungal and mycoplasma contaminants, and stocks of each were stored frozen in liquid nitrogen. Cultures were grown as monolayers or in suspension in standard cell culture media. Cells were recovered by scraping (from monolayers) or by centrifugation (from suspension cultures), and were washed two or three times by suspension in phosphate buffered saline followed by collection by centrifugation. Cell pellets were stored frozen at -40°C. Crude cell membranes were prepared by homogenisation followed by high-speed centrifugation, and characterisation of cloned receptors achieved by radioligand binding.

Preparation of CHO cell membranes

Cell pellets were gently thawed at room temperature, and resuspended in about 20 volumes of ice-cold 50 mM Tris salts (pH 7.4 @ 37°C), 20mM EDTA, 0.2 M sucrose. The suspension was homogenised using an Ultra-Turrax at full speed for 15 sec. The homogenate was centrifuged at 18,000 r.p.m for 20 min at 4°C in a Sorvall RC5C centrifuge. The membrane pellet was resuspended in ice-cold 50 mM Tris salts (pH 7.4 @ 37°C), using an Ultra-Turrax, and recentrifuged at 18,000 r.p.m for 15 min at 4°C in a Sorvall RC5C. The membranes were washed two more times with ice-cold 50 mM Tris salts (pH 7.4 @ 37°C). The final pellet was resuspended in 50 mM Tris salts (pH 7.4 @ 37°C), and the protein content determined using bovine serum albumin as a standard (Bradford, M. M. (1976) Anal. Biochem. 72, 248-254).

Binding experiments on cloned dopamine receptors

Crude cell membranes were incubated with 0.1 nM [¹²⁵I] idosulpride (~2000 Ci/mmol; Amersham, U. K.), and the test compound in a buffer containing 50 mM Tris salts (pH 7.4 @ 37°C), 120 mM NaCl, 5 mM KCl, 2 mM CaCl₂, 1 mM MgCl₂, 0.1% (w/v) bovine serum albumin, in a total volume of 1 ml for 30 min at 37°C. Following incubation, samples were filtered using a Brandel Cell Harvester, and washed three times with ice-cold 50 mM Tris salts (pH 7.4 @ 37°C), 120 mM NaCl, 5 mM KCl, 2 mM CaCl₂, 1 mM MgCl₂. The radioactivity on the filters was measured using a Cobra gamma counter (Canberra Packard). Non-specific binding was defined as the radioligand binding remaining after incubation in the presence of 100 µM idosulpride. For competition curves, 14 concentrations (half-log dilutions) of competing cold drug were used. Competition curves were analysed simultaneously whenever possible using non-linear least-squares fitting procedures, capable of fitting one, two or three site models.

Compounds of Examples tested according to this method had pK_i values in the range 7.0 - 9.0 at the human cloned dopamine D₃ receptor.

Functional Activity at cloned dopamine receptors

The functional activity of compounds at human D2 and human D3 receptors (ie agonism or antagonism) may be determined using a Cytosensor Microphysiometer (McConnell HM et al Science 1992 257 1906-1912) In Microphysiometer experiments, cells

5 (hD2_CHO or hD3_CHO) were seeded into 12mm Transwell inserts (Costar) at 300000 cells/cup in foetal calf serum (FCS)-containing medium. The cells were incubated for 6h at 37°C in 5%CO₂, before changing to FCS-free medium. After a further 16-18h, cups were loaded into the sensor chambers of the Cytosensor Microphysiometer (Molecular

10 Devices) and the chambers perfused with running medium (bicarbonate-free Dulbecco's modified Eagles medium containing 2 mM glutamine and 44 mM NaCl) at a flow rate of 100 ul/min. Each pump cycle lasted 90s. The pump was on for the first 60s and the acidification rate determined between 68 and 88s, using the Cytosoft programme. Test compounds were diluted in running medium. In experiments to determine agonist

15 activity, cells were exposed (4.5 min for hD2, 7.5 min for hD3) to increasing concentrations of putative agonist at half hour intervals. Seven concentrations of the putative agonist were used. Peak acidification rate to each putative agonist concentration was determined and concentration-response curves fitted using Robofit [Tilford, N.S., Bowen, W.P. & Baxter, G.S. Br. J. Pharmacol. (1995) in press]. In experiments to

20 determine antagonist potency, cells were treated at 30 min intervals with five pulses of a submaximal concentration of quinpirole (100 nM for hD2 cells, 30 nM for hD3 cells), before exposure to the lowest concentration of putative antagonist. At the end of the next 30 min interval, cells were pulsed again with quinpirole (in the continued presence of the antagonist) before exposure to the next highest antagonist concentration. In all, five

25 concentrations of antagonist were used in each experiment. Peak acidification rate to each agonist concentration was determined and concentration-inhibition curves fitted using Robofit.

Compounds of Examples tested according to this method were shown to be antagonists with pK_b values in the range 7.0 - 10.0 at the human cloned dopamine D₃ receptor.

30

Pharmaceutical Formulations

The following represent typical pharmaceutical formulations according to the present invention, which may be prepared using standard methods.

35 IV Infusion

Compound of formula (I)	1-40 mg
Buffer	to pH ca 7
Solvent/complexing agent	to 100 ml

Bolus Injection

40 Compound of formula (I)	1-40 mg
Buffer	to pH ca 7
Co-Solvent	to 5 ml

Buffer : Suitable buffers include citrate, phosphate, sodium hydroxide/hydrochloric acid.

5 Solvent : Typically water but may also include cyclodextrins (1-100 mg) and co-solvents such as propylene glycol, polyethylene glycol and alcohol.

Tablet

Compound	1 - 40 mg
Diluent/Filler *	50 - 250 mg
Binder	5 - 25 mg
10 Disintegrant *	5 - 50 mg
Lubricant	1 - 5 mg
Cyclodextrin	1 - 100 mg

* may also include cyclodextrins

15

Diluent : e.g. Microcrystalline cellulose, lactose, starch

Binder : e.g. Polyvinylpyrrolidone, hydroxypropymethylcellulose

Disintegrant : e.g. Sodium starch glycollate, crospovidone

Lubricant : e.g. Magnesium stearate, sodium stearyl fumarate.

20

Oral Suspension

Compound	1 - 40 mg
Suspending Agent	0.1 - 10 mg
Diluent	20 - 60 mg
25 Preservative	0.01 - 1.0 mg
Buffer	to pH ca 5 - 8
Co-solvent	0 - 40 mg
Flavour	0.01 - 1.0 mg
Colourant	0.001 - 0.1 mg

30

Suspending agent : e.g. Xanthan gum, microcrystalline cellulose

Diluent : e.g. sorbitol solution, typically water

Preservative : e.g. sodium benzoate

Buffer : e.g. citrate

35 Co-solvent : e.g. alcohol, propylene glycol, polyethylene glycol, cyclodextrin

The invention is further illustrated by the following non-limiting examples :

Description 1

40

2,3,4,5-Tetrahydro-1H-3-benzazepine

1,2-Phenylenediacetonitrile (7.5g, 48 mmol) dissolved in ethanol (150ml) was added to Raney Ni (2g) which had been previously washed with ethanol (3x20ml). The mixture was then hydrogenated at 50°C at 50psi pressure with shaking for 24h. The reaction mixture was then cooled to room temperature and filtered through a pad of kieselguhr and washed through with ethanol (100ml). The filtrate was evaporated *in vacuo* to give a brown oil which was chromatographed on silica gel (100g), eluting with 2-10% methanol in CH₂Cl₂ to give the title compound as a brown oil (2.45g, 35%).

Mass spectrum (API⁺) Found: 148 (MH⁺). C₁₀H₁₃N requires 147.

Description 2

trans-3-(2-(1-(4-(N-tert-Butoxycarbonyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine

Sodium triacetoxyborohydride (4.3g, 20.4 mmol) was added to a mixture of 2,3,4,5-tetrahydro-1H-3-benzazepine (2.0g, 13.6 mmol), and *trans-2-(1-(4-(N-tert-butoxycarbonyl)amino)cyclohexyl)acetaldehyde* in 1,2-dichloroethane (200ml), and the mixture stirred at room temperature for 0.5h. The reaction mixture was diluted with CH₂Cl₂ (100ml) and washed with saturated aqueous K₂CO₃ (200ml), followed by brine (100ml). The organic layer was separated and dried over Na₂SO₄, then evaporated *in vacuo* to give an off-white solid which was chromatographed on silica gel eluting with ethyl acetate to give the title compound as an off-white solid (3.13g, 62%).

Mass spectrum (API⁺): Found 373. C₂₃H₃₆N₂O₂ requires 372.

The following compound was prepared in a similar manner to Description 2

(a) *trans-3-(2-(1-(4-N-tert-Butoxycarbonyl)amino)cyclohexyl)ethyl-6-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine*

Mass spectrum (API⁺): Found 403 (MH⁺). C₂₄H₃₈N₂O₃ requires 402.

b) *trans-3-(2-(1-(4-N-tert-Butyloxycarbonyl)amino)cyclohexyl)ethyl-7-cyano-2,3,4,5-tetrahydro-1H-3-benzazepine*

Mass spectrum (API⁺) Found 398 (MH⁺). C₂₄H₃₅N₃O₂ requires 397.

¹H NMR (CDCl₃) δ: 0.97 – 1.13 (4H, m), 1.22 (1H, m), 1.36 – 1.47 (11H, m), 1.71 – 1.79 (2H, m), 1.95 – 2.04 (2H, m), 2.48 (2H, m), 2.61 (4H, m), 2.90 – 3.00 (4H, m), 3.37 (1H, m), 4.35 (1H, m), 7.17 (1H, d, J = 5 Hz), 7.36 (1H, s), 7.52 (1H, d, J = 5 Hz).

Description 3**trans-3-(2-(1-(4-Amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine**

A mixture of *trans*-2-(2-(1-(4-(*N*-*tert*-butoxycarbonyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine (3.1g, 8.3 mmol) and trifluoroacetic acid (5ml) in CH₂Cl₂ (50ml) was stirred at room temperature for 1h, then at 40°C for 1h. The reaction mixture was then diluted with CH₂Cl₂ (100ml) and washed with saturated aqueous K₂CO₃ (2x100ml). The organic layer was dried over Na₂SO₄ and evaporated *in vacuo* to give the title compound as a brown oil (2.14g, 95%).

Mass spectrum (API⁺): Found 273 (MH⁺). C₁₈H₂₈N₂ requires 272.

The following compound was prepared in a similar manner to Description 3

(a) trans-3-(2-(1-(4-Amino)cyclohexyl)ethyl)-6-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine

Mass spectrum (API⁺): Found 303 (MH⁺). C₁₉H₃₀N₂O requires 302.

b) trans-3-(2-(1-(4-Amino)cyclohexyl)ethyl)-7-cyano-2,3,4,5-tetrahydro-1H-3-benzazepine

Mass spectrum (API⁺): Found 298 (MH⁺). C₁₉H₂₇N₃ requires 297.

¹H NMR (CDCl₃) δ: 0.92 – 1.18 (6H, m), 1.21 (1H, m), 1.41 (2H, m), 1.75 (2H, m), 1.85 (2H, m), 2.49 (2H, m), 2.60 (5H, m), 2.95 (4H, m), 7.16 (1H, d, J = 5 Hz), 7.36 (1H, s), 7.40 (1H, d, J = 5 Hz).

Description 4**trans-2-(1-(4-(N-tert-Butyloxycarbonyl)amino)cyclohexyl)acetic acid, methyl ester**

A mixture of *trans*-(4-amino)cyclohexylactic acid hydrogen sulfate (T.P. Johnston *et al*; J. Med Chem., 1977, 20 (2), 279-290), (27.0g, 106mmol), conc. H₂SO₄ (3ml), and methanol (300ml) was stirred at reflux for 5h. Resulting solution was filtered and the filtrate evaporated *in vacuo* to give a brown oil (36g). A mixture of this material, triethylamine (36ml; 26.1g, 259 mmol), dichloromethane (600ml) and di-*t*-butyl dicarbonate (25.5g, 117mmol) was stirred at 20°C for 18h. Resulting solution was partitioned between saturated aqueous NaHCO₃ (500ml) and dichloromethane (3x200ml), and the combined extracts were dried (Na₂SO₄) and evaporated *in vacuo* to give the title compound (24.6g, 86%) as a colourless solid.

¹H NMR (CDCl₃) δ: 1.08 (4H, m), 1.43 (9H, s), 1.76 (3H, m), 2.00 (2H, m), 2.20 (2H, d, J = 7 Hz), 3.37 (1H, m), 3.66 (3H, s), 4.39 (1H, br s).

5 Description 5

trans-2-(1-(4-(*N*-*tert*-Butyloxycarbonyl)amino)cyclohexyl)acetaldehyde

To a stirred solution of *trans*-2-(1-(4-(*N*-*tert*-butyloxycarbonyl)amino)cyclohexyl)acetic acid, methyl ester (46.0g, 170 mmol) in dry toluene (920ml) at -78°C under argon was added a solution of di-isobutylaluminium hydride (1M; 285 ml; 285 mmol), dropwise over 0.5h. Resulting solution was stirred for a further 0.3h and quenched with a mixture of methanol (28ml) in toluene (50ml) and then poured into saturated aqueous potassium sodium tartrate (1.2L). The resultant mixture was extracted with ether (4x1L). The combined organic extracts were dried (Na₂SO₄) and evaporated *in vacuo* to give a waxy solid which was purified using silica gel, eluting with 10-50% ethyl acetate/hexane to give the title compound (21.77g, 53%) as a colourless solid.

¹H NMR (CDCl₃) δ: 1.12 (4H, m), 1.44 (9H, s), 1.78 (3H, m), 2.00 (2H, m), 2.33 (2H, dd, J = 7, 2 Hz), 3.37 (1H, m), 4.40 (1H, m), 9.75 (1H, m).

Description 6

7-Hydroxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine, hydrobromide

7-Methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine (10 g) in 48% aqueous hydrobromic acid (350 ml) was allowed to stir at 100 °C for 4 h. The mixture was cooled to 20 °C then evaporated to dryness *in vacuo* to give the title compound (14.5 g) as a brown solid.

Mass spectrum (API⁺): Found 164 (MH⁺). C₁₀H₁₃NO requires 163.

¹H NMR (DMSO) δ: 2.80 – 3.25 (8H, m), 4.42 (2H, br s), 6.50 – 6.70 (2H, m), 6.98 (1H, d, J = 8 Hz), 8.86 (1H, br s).

35 Description 7

3-(*tert*-Butyloxycarbonyl)-7-hydroxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine

To a solution of 7-hydroxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine, hydrobromide (14.5 g) in tetrahydrofuran (100 ml) and water (70 ml), was added triethylamine (8 g), followed by a solution of di-*tert*-butyl dicarbonate (14 g) in THF (20 ml). The resulting mixture was allowed to stir at 20 °C for 16 h, partitioned between ethyl acetate (200 ml) and water (200 ml). The aqueous layer was washed with ethyl acetate (100 ml). The combined organic extracts were washed with saturated aqueous sodium bicarbonate (100 ml), dried

(Na₂SO₄) and evaporated to dryness *in vacuo*. The resulting oil was purified by silica gel chromatography. Elution with ethyl acetate in hexane (10% - 30%) gave the title compound (8 g).

- 5 Mass spectrum (API⁺): Found 164 (MH⁺-Boc). C₁₅H₂₁NO₃ requires 263.

¹H NMR (CDCl₃) δ: 1.48 (9H, s), 2.75 – 2.87 (4H, m), 3.40 – 3.60 (4H, m), 4.95 (1H, s), 6.50 – 6.62 (2H, m), 6.96 (1H, d, J = 8 Hz).

10 **Description 8**

3-(*tert*-Butyloxycarbonyl)-7-trifluoromethylsulfonyloxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine

- 15 To a stirred mixture of 3-(*tert*-butyloxycarbonyl)-7-hydroxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine (7 g) and triethylamine (5.4 ml) in dry dichloromethane under argon at -20 °C, was added, dropwise, trifluoromethanesulfonic anhydride (5 ml). The resulting mixture was allowed to warm slowly to 20 °C over 16 h, then was poured into saturated aqueous sodium bicarbonate (200 ml) and extracted with dichloromethane (2 x 150 ml).
20 The combined organic extracts were washed with brine (150 ml), dried (Na₂SO₄) and evaporated *in vacuo* to give an amber oil. Silica gel chromatography, eluting with ethyl acetate in hexane (10% - 30%) gave the title compound (7 g) as an amber oil.

- 25 Mass spectrum (API⁺): Found 396 (MH⁺). C₁₆H₂₀F₃NO₅S requires 395.

¹H NMR (CDCl₃) δ: 1.48 (9H, s), 2.85 – 2.95 (4H, m), 3.5 – 3.65 (4H, m), 7.00 – 7.05 (2H, m), 7.15 – 7.27 (1H, m).

Description 9

30

3-(*tert*-Butyloxycarbonyl)-7-cyano-2,3,4,5-tetrahydro-1*H*-3-benzazepine

- A mixture of 3-(*tert*-butoxycarbonyl)-7-trifluoromethylsulfonyloxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine (4.78 g, 12.1 mmol), zinc cyanide (1.42 g, 15.6 mmol) and *tetrakis*-
35 triphenylphosphine palladium (0) (1.4 g, 1.2 mmol, 10 mol%), in dry dimethylformamide (50ml) was stirred at 100 °C for 3 h under argon. After cooling to room temperature the reaction mixture was diluted with ethyl acetate (120 ml) and filtered. The filtrate was washed with saturated aqueous sodium bicarbonate (100 ml), then water (2 x 50 ml), then brine (50 ml). The organic layer was dried over sodium sulfate and evaporated *in vacuo*
40 to give brown oil, which was purified by chromatography on silica gel with 20 – 100% ethyl acetate - hexane elution to give the title compound (0.765 g, 23%) as a brown oil.

Mass spectrum (API⁺): Found 173 (MH⁺-Boc). C₁₆H₂₀N₂O₂ requires 272.

¹H NMR (CDCl₃) δ: 1.47 (9H, s), 2.93 (4H, m), 3.56 (4H, m), 7.21 (1H, d, J = 8 Hz), 7.42 (2H, m).

Description 10

5

7-Cyano-2,3,4,5-tetrahydro-1H-3-benzazepine

A mixture of 3-(*tert*-butoxycarbonyl)-7-cyano-2,3,4,5-tetrahydro-1H-3-benzazepine (765 mg, 2.81 mmol) and trifluoroacetic acid (2 ml), in dichloromethane (20 ml) was stirred at 40 °C for 1 h. The reaction mixture was evaporated to dryness *in vacuo* and partitioned between ethyl acetate (50 ml) and water (50 ml). The aqueous layer was basified using potassium carbonate and re-extracted with ethyl acetate (2 x 30ml). The combined basic organic extracts were dried over sodium sulfate and evaporated *in vacuo* to give the title compound as a colourless oil (212 mg, 44%).

15

Mass spectrum (API⁺): Found 173 (MH⁺). C₁₁H₁₂N₂ requires 172.

¹H NMR (CDCl₃) δ: 2.04 (1H, br s), 2.95 (8H, m), 7.18 (1H, d, J = 8 Hz), 7.38 (2H, m).

20 Description 11

7-Acetyl-*trans*-3-(2-(1-(4-*N*-*tert*-butoxycarbonyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine

25 Mass spectrum (API⁺): Found 415 (MH⁺). C₂₅H₃₈N₂O₃ requires 414.

Description 12

30 7-Acetyl-*trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine

Mass spectrum (API⁺): Found 315 (MH⁺). C₂₀H₃₀N₂O requires 314.

35 ¹H NMR (CDCl₃) δ: 0.80 - 1.30 (5H, m), 1.41 (4H, m), 1.65 - 1.85 (4H, m), 2.48 (2H, m), 2.57 (3H, s), 2.60 (5H, m), 2.97 (4H, m), 7.17 (1H, d, J = 8 Hz), 7.70 (2H, m).

Description 13

40 7-Acetyl-3-(*tert*-butyloxycarbonyl)-2,3,4,5-tetrahydro-1H-3-benzazepine

To a stirred solution of 3-(*tert*-butyloxycarbonyl)-7-trifluoromethylsulfonyloxy-2,3,4,5-tetrahydro-1H-3-benzazepine (10 g, 25.3 mmol) in anhydrous dimethylformamide (100 ml) under argon at room temperature, was added triethylamine (7.05 ml, 50.6 mmol), butyl vinyl ether (16.4 ml, 126.6 mmol), 1,3-bis(diphenylphosphino)propane (0.412 g, 1

mmol) and palladium acetate (0.202 g, 0.9 mmol) sequentially. The resultant mixture was heated at 85°C for 1.5 h and cooled to room temperature. 4 % Aqueous hydrochloric acid (150 ml) was added and stirring continued for 0.5 h. The reaction mixture was extracted with dichloromethane (3 x 300 ml) and the combined organics washed with water (4 x 500 ml), dried (Na₂SO₄) and evaporated *in vacuo* to afford a brown gum. Chromatography on silica gel with 0 - 30% ethyl acetate - hexane gradient elution gave the title compound (5.8 g, 79 %) as a colourless solid.

¹H NMR (CDCl₃) δ: 1.49 (9H, s), 2.58 (3H, s), 2.96 (4H, m), 3.57 (4H, m), 7.21 (1H, d, J = 8 Hz), 7.72 (2H, m).

Description 14

7-Acetyl-2,3,4,5-tetrahydro-1H-benzazepine

Prepared from 7-acetyl-3-(*tert*-butoxycarbonyl)-2,3,4,5-tetrahydro-1H-3-benzazepine (6.3 g, 21.8 mmol) using the method of Description 10 to afford a pale yellow oil (4.12 g, 100 %).

¹H NMR (CDCl₃) δ: 1.89 (1H, s), 2.58 (3H, s), 2.97 (8H, s), 7.17 (1H, d, J = 8 Hz), 7.70 (2H, m).

Description 15

3-(3-Bromophenyl)-5-methyl-1,2,4-oxadiazole

Potassium *tert*-butoxide (7.33 g, 65.4 mmol) was added over 5 minutes to ice chilled, stirred methanol under argon. After a further 5 min hydroxylamine hydrochloride (4.9 g, 70.43 mmol) was added in one portion and the resultant mixture stirred at room temperature for 1 h. A solution of 3-bromobenzonitrile (7.93 g, 43.6 mmol) in methanol (120 ml) was added in one portion and the mixture heated at reflux for 4 h, cooled filtered, and the filtrate evaporated *in vacuo*. The residue was refluxed in acetic anhydride (60 ml) for 3 h, cooled to room temperature and poured into ice-water (300 ml). The precipitate was filtered, washed with water, dried *in vacuo* and chromatographed on silica eluting with 0 - 10% ethyl acetate - hexane gradient. Fractions containing desired product were pooled and evaporated *in vacuo* and the residue recrystallised from hexane to afford the title compound as colourless crystals (5.2 g, 50 %).

Mass spectrum: (API⁺) Found: 239 (MH⁺). C₉H₇⁷⁹BrN₂O requires 238

¹H NMR (CDCl₃) δ: 2.66 (3H, s), 7.36 (1H, t, J = 8 Hz), 7.63 (1H, m), 8.05 (1H, m), 8.23 (1H, m).

Description 16**3-(5-Methyl-1,2,4-oxadiazol-3-yl)-benzoic acid**

- 5 A mixture of 3-(3-bromophenyl)-5-methyl-1,2,4-oxadiazole (2.68 g, 11.3 mmol), tributylamine (3.05 ml, 12.5 mmol) and *trans*-dibromobis(triphenylphosphine)palladium (II) (0.13 g, 0.16 mmol) in methanol (5 ml) was carbonylated at 30 psi and 100 °C for 18 h. The mixture was cooled to room temperature, diluted with ethyl acetate (100 ml) and washed sequentially with saturated sodium hydrogen carbonate (2 x 300 ml), brine (100
10 ml), 0.5 N hydrochloric acid (200 ml), brine (100 ml), then dried Na₂SO₄ and evaporated *in vacuo* to afford a yellow oil (2.49 g). A 2 g sample of this oil was dissolved in aqueous methanol (5:3, 80 ml), sodium hydroxide (0.36 g) added and the mixture stirred at room temperature for 20 h. The mixture was evaporated *in vacuo* and the residue partitioned between ethyl acetate (100 ml) and water (100 ml). The aqueous
15 layer was acidified with 2N HCl and the resultant precipitate filtered, washed with water and dried *in vacuo* to afford the title compound as a colourless solid (0.78 g, 42 %).

Mass spectrum: (API⁺) Found: 205 (MH⁺). C₁₀H₈N₂O₃ requires 204.

- 20 ¹H NMR (CDCl₃) δ: 2.70 (3H, s), 7.71 (1H, m), 8.14 (1H, dd, J = 7, 1 Hz), 8.23 (1H, dd, J = 7, 1 Hz), 8.54 (1H, m), 13.35 (1H, br s).

Description 17**3-(1-Pyrazolyl)-benzoic acid**

- A mixture of 3-hydrazinobenzoic acid (1.52 g, 0.01 mmol) and malondialdehydebis(dimethylacetal) (2.39 ml; 0.01 mol) in ethanol (10 ml) and water (15 ml) was heated at reflux for 2 h. The resulting solution was cooled and evaporated to
30 afford the title product (1.8 g,; 96 %) as a yellow solid.

- ¹H NMR (DMSO-d₆) δ: 6.60 (1H, t, J = 2 Hz), 7.65 (1H, t, J = 8 Hz), 7.81 (1H, d, J = 1.5 Hz), 7.89 (1H, dd, J = 8 and 1.5 Hz), 8.12 (1H, dd, J = 8 and 1.5 Hz), 8.4 (1H, d, J = 2 Hz), 8.64 (1H, d, J = 2 Hz).

35

Mass spectrum (API⁺): Found 189 (MH⁺). C₁₀H₈N₂O₂ requires 188.

Description 18

- 40 **3-(*tert*-Butoxycarbonyl)-7-(3-(5-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1H-3-benzazepine**

- To a suspension of sodium methoxide (0.6 g, 11 mmol) in anhydrous methanol (12ml) under argon, was added hydroxylamine hydrochloride (0.76 g, 11 mmol), followed by 3-(*tert*-butoxycarbonyl)-7-cyano-2,3,4,5-tetrahydro-1*H*-3-benzazepine (1.5 g, 5.5 mmol). The mixture was stirred under reflux for 16 h, then allowed to cool to room temperature.
- 5 The methanol was evaporated *in vacuo* and the resulting residue partitioned between dichloromethane (100 ml) and water (100 ml). The aqueous layer was washed with more CH₂Cl₂ (100 ml). The combined organic extracts were dried and evaporated *in vacuo* to give a solid (1.8 g), which was mixed with acetic anhydride (15 ml) and heated at 120 °C for 2 h. Excess acetic anhydride was evaporated *in vacuo* and the resulting oily residue
- 10 partitioned between CH₂Cl₂ (250 ml) and saturated sodium bicarbonate solution (250 ml). The organic layer was washed with more bicarbonate solution (200 ml), dried, and evaporated to give an oil. Gravity silica gel chromatography eluting with ethyl acetate in hexane gave the title compound (3.2 g, 73 %) as a colourless oil.
- 15 ¹H NMR (CDCl₃) δ: 1.49 (9H, s), 2.65 (3H, s), 2.96 (4H, m), 3.58 (4H, m), 7.22 (1H, d, J = 8 Hz), 7.80 (2H, m).

Description 19

20 **7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

- A solution of 3-(*tert*-butoxycarbonyl)-7-(3-(5-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine (1.2 g, 3.6 mmol) in CH₂Cl₂ (15 ml) and trifluoroacetic acid (15 ml) was heated under reflux for 2 h. Solvent was evaporated *in vacuo* and the residue
- 25 partitioned between diethyl ether (50 ml) and water (50 ml). The aqueous layer was saturated with potassium carbonate then extracted with CH₂Cl₂ (2 x 100 ml). The combined organic extracts were dried and evaporated *in vacuo* to give the title compound (0.74 g, 88 %) as an oil.
- 30 Mass spectrum (API⁺): Found 230 (MH⁺). C₁₃H₁₃N₃O requires 229.
- ¹H NMR (CDCl₃) δ: 1.80 (1H, br s), 2.65 (3H, s), 2.90 - 3.00 (8H, m), 7.20 (1H, d, J = 8 Hz), 7.75 - 7.85 (2H, m).

Description 20**7-(3-(*tert*-Butoxycarbonyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepinyl)carboxamide**

- 5 To a solution of 3-(*tert*-butoxycarbonyl)-7-cyano-2,3,4,5-tetrahydro-1*H*-3-benzazepine (5.44 g, 20 mmol) cooled in ice bath, was added potassium carbonate (0.4 g) in water (1 ml), followed by dropwise addition of 30 % w/w hydrogen peroxide (2.4 ml). The resulting mixture was stirred at 5 °C for 5 min, then the ice-bath was removed. After another 5 min, water (100 ml) was added. The solid precipitate was collected by filtration and dried to give the title compound (4.35 g, 75 %) as a colourless solid.

¹H NMR (CDCl₃) δ: 1.48 (9H, s), 2.96 (4H, m), 3.56 (4H, m), 5.60 - 6.30 (2H, br d), 7.19 (1H, d, J = 8 Hz), 7.50 - 7.80 (2H, m).

15 Description 21**3-(*tert*-Butoxycarbonyl)-7-(5-(3-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

- 20 A mixture of 7-(3-(*tert*-butoxycarbonyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepinyl)carboxamide (4.29 g, 14.8 mmol) and *N,N*-dimethyl acetamide dimethyl acetal (6 ml, 41 mmol) was heated at 125 °C under argon. Methanol was removed from the reaction by means of a distillation condenser over 2 h. The reaction mixture was further evaporated *in vacuo* to give a thick brown oily residue. To this residue was added, in order, dioxan (10 ml), 5M sodium hydroxide (4 ml), hydroxylamine hydrochloride (1.4 g, 20 mmol) and 70 % aqueous acetic acid (20 ml). The combined mixture was allowed to stir at room temperature for 15 min and then at 90 °C for 1h. The mixture was treated with water (100 ml) and extracted with CH₂Cl₂ (2x150 ml). Combined organic extracts were washed with saturated sodium bicarbonate (100 ml), dried and evaporated *in vacuo* to give an oil. Gravity silica gel chromatography, eluting with ethyl acetate in hexane, gave the title compound (3.9 g, 80 %) as a colourless solid.

¹H NMR (CDCl₃) δ: 1.49 (9H, s), 2.47 (3H, s), 2.98 (4H, m), 3.60 (4H, m), 7.27 (1H, d, J = 8 Hz), 7.80 - 7.90 (2H, m).

35 Description 22**7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

- 40 A solution of 3-(*tert*-butoxycarbonyl)-7-(5-(3-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine (3.8 g, 11.6 mmol) in CH₂Cl₂ (50 ml) and trifluoroacetic acid (12 ml) was heated under reflux for 2 h. Solvent was evaporated *in vacuo* and the residue partitioned between diethyl ether (200 ml) and water (200 ml). The aqueous layer was saturated with potassium carbonate then extracted with CH₂Cl₂ (3 x 200 ml). The

combined organic extracts were dried and evaporated *in vacuo* to give the title compound (2.4 g, 91 %) as a colourless solid.

Mass spectrum (API⁺): Found 230 (MH⁺). C₁₃H₁₅N₃O requires 229.

¹H NMR (CDCl₃) δ: 1.86 (1H, br s), 2.47 (3H, s), 3.00 (8H, m), 7.25 (1H, d, J = 8 Hz), 7.80 - 7.90 (2H, m).

Description 23

***trans*-3-(2-(1-(4-*N*-*tert*-Butyloxycarbonyl)amino)cyclohexyl)ethyl-7-(3-(5-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

Prepared from 7-(3-(5-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine in a manner similar to Description 2, in 96 % yield.

Mass spectrum (API⁺): Found 455 (MH⁺). C₂₆H₃₈N₄O₃ requires 454.

¹H NMR (CDCl₃) δ: 0.90 - 1.10 (4H, m), 1.15 - 1.25 (1H, m), 1.38 - 1.47 (11H, m), 1.73 - 1.85 (2H, m), 1.93 - 2.05 (2H, m), 2.40 - 2.55 (2H, m), 2.56 - 2.70 (7H, m), 2.90 - 3.05 (4H, m), 3.35 (1H, br s), 4.35 (1H, br s), 7.19 (1H, d, J = 8 Hz), 7.75 - 7.85 (2H, m).

Description 24

***trans*-3-(2-(1-(4-*N*-*tert*-Butyloxycarbonyl)amino)cyclohexyl)ethyl-7-(5-(3-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

Prepared from 7-(5-(3-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine in a manner similar to Description 2, in 94 % yield.

Mass spectrum (API⁺): Found 455 (MH⁺). C₂₆H₃₈N₄O₃ requires 454.

¹H NMR (CDCl₃) δ: 0.95 - 1.10 (4H, m), 1.23 (1H, br s), 1.40 - 1.50 (11H, m), 1.70 - 1.85 (2H, m), 1.95 - 2.10 (2H, m), 2.46 (3H, s), 2.46 - 2.52 (2H, m), 2.60 - 2.70 (4H, m), 2.90 - 3.60 (4H, m), 3.35 (1H, m), 4.35 (1H, m), 7.23 (1H, d, J = 8 Hz), 7.80 - 7.90 (2H, m).

Description 25

***trans*-3-(2-(1-(4-Amino)cyclohexyl)ethyl)-7-(3-(5-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

Prepared from *trans*-3-(2-(1-(4-*N*-*tert*-butyloxycarbonyl)amino)cyclohexyl)ethyl-7-(3-(5-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine in a manner similar to Description 3, in 100 % yield.

Mass spectrum (API⁺): found 355 (MH⁺). C₂₁H₃₀N₄O requires 354.

5 ¹H NMR (CDCl₃) δ: 0.90 - 1.10 (4H, m), 1.40 (2H, br s), 1.12 - 1.25 (1H, m), 1.40 - 1.50 (2H, m), 1.70 - 1.80 (2H, m), 1.80 - 1.90 (2H, m), 2.40 - 2.50 (2H, m), 2.55 - 2.70 (8H, m), 2.90 - 3.00 (4H, m), 7.19 (1H, d, J = 8 Hz), 7.75 - 7.85 (2H, m).

Description 26

10 ***trans*-3-(2-(1-(4-Amino)cyclohexyl)ethyl)-7-(5-(3-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1H-3-benzazepine**

15 Prepared from *trans*-3-(2-(1-(4-*N*-*tert*-butoxycarbonyl)amino)cyclohexyl)ethyl-7-(5-(3-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1H-3-benzazepine in a manner similar to Description 3, in 100 % yield.

Mass spectrum (API⁺): Found 355 (MH⁺). C₂₁H₃₀N₄O requires 354.

20 ¹H NMR (CDCl₃) δ: 0.90 - 1.30 (5H, m), 1.37 - 1.50 (2H, m), 1.64 (2H, br s), 1.70 - 1.95 (4H, m), 2.46 (3H, s), 2.46 - 2.70 (7H, m), 2.90 - 3.10 (4H, m), 7.24 (1H, d, J = 8 Hz), 7.80 - 7.90 (2H, m).

Description 27

25 **3-Acetyl-2,3,4,5-tetrahydro-1H-3-benzazepine**

A solution of acetic anhydride (6.37 g, 0.062 mol) in dichloromethane (50 ml) was added dropwise to a stirred solution of 2,3,4,5-tetrahydro-1H-3-benzazepine (8.35 g, 0.057 mol) and triethylamine (8.7 ml) in dichloromethane (50 ml) at 0 °C under argon. After stirring at room temperature for 18 h, water (80 ml) was added and the organic layer separated.

30 The organic layer was washed with 0.5 M hydrochloric acid (50 ml), saturated sodium bicarbonate solution (50 ml), water (50 ml) and then dried (Na₂SO₄). Evaporation of the solvent *in vacuo* gave the title compound (10.24 g, 95 %) as a yellow oil which solidified on standing.

35 ¹H NMR (CDCl₃) δ: 2.18 (3H, s), 2.85 - 3.00 (4H, m), 3.55 - 3.60 (2H, m), 3.72 - 3.80 (2H, m), 7.10 - 7.20 (4H, m).

Mass Spectrum AP⁺: Found 190 (MH⁺). C₁₂H₁₅NO requires 189.

Description 28

3-Acetyl-7-chlorosulphonyl-2,3,4,5-tetrahydro-1H-3-benzazepine

A solution of 3-acetyl-2,3,4,5-tetrahydro-1H-3-benzazepine (4.0 g, 0.021 mol) in dichloromethane (25 ml) was added dropwise to a stirred solution of chlorosulphonic acid in dichloromethane (25 ml) at -70 °C under argon. After warming to room temperature, the reaction was stirred for 18 h before being quenched in ice/water (200 ml). The resulting mixture was extracted with ethyl acetate (3 x 100 ml), dried (Na₂SO₄) and the solvent evaporated *in vacuo* to give the title compound (2.74 g, 45 %) as a pale yellow solid.

¹H NMR: δ (CDCl₃): 2.21 (3H, s), 3.0 - 3.10 (4H, m), 3.60 - 3.70 (2H, m), 3.74 - 3.80 (2H, m), 7.35 - 7.40 (1H, m), 7.80 - 7.85 (2H, m).

Mass spectrum AP⁺: Found 288 & 290 (MH⁺). C₁₂H₁₄NSO₂Cl requires 287 & 289.

Description 29**3-Acetyl-7-methylsulphonyl-2,3,4,5-tetrahydro-1H-3-benzazepine**

To a stirred solution of sodium sulphite (1.60 g, 12.6 mmol) and sodium hydrogen carbonate (1.14 g, 13.56 mmol) in water (25 ml) was added 7-3-acetyl-7-chlorosulphonyl-2,3,4,5-tetrahydro-1H-3-benzazepine (2.6 g, 9.04 mmol) in tetrahydrofuran (10 ml). The reaction mixture was then heated at 75 °C for 2 h, cooled to 30 °C and methyl iodide (2.8 ml, 45.20 mmol) added. After stirring at 50 °C for 24 h, the reaction mixture was cooled to room temperature and partitioned between water (50 ml) and ethyl acetate (100 ml). The aqueous layer was then separated and further extracted with ethyl acetate (2 x 80 ml). The combined organics were then dried (Na₂SO₄) and the solvent removed *in vacuo* to give the title compound (1.77 g, 73 %) as a pale yellow solid.

¹H NMR (CDCl₃) 2.20 (3H, s), 2.99 - 3.05 (4H, m), 3.06 (3H, s), 3.61 - 3.64 (2H, m), 3.73 - 3.77 (2H, m), 7.32 - 7.37 (1H, m), 7.7 - 7.75 (2H, m).

Mass Spectrum AP⁺: Found 268 (MH⁺). C₁₃H₁₇NSO₃ requires 267.

Description 30**7-Methylsulphonyl-2,3,4,5-tetrahydro-1H-3-benzazepine**

A solution of 3-acetyl-7-methylsulphonyl-2,3,4,5-tetrahydro-1H-3-benzazepine (1.75 g, 6.55 mmol) in 5 M hydrochloric acid was heated at reflux for 18 h. The reaction mixture was then cooled to room temperature, basified to pH = 12 with potassium carbonate and the solvent evaporated *in vacuo*. The solid residue was then extracted with ethyl acetate

(5 x 60 ml) and the combined organics dried (Na_2SO_4). The solvent was then evaporated *in vacuo* to give the title compound (450 mg, 32 %) as a pale yellow oil.

^1H NMR (CDCl_3) 1.88 (1H, br s), 2.95 - 3.13 (8H, m), 3.04 (3H, s), 7.25 - 7.30 (1H, d),
5 7.65 - 7.72 (2H, m).

Mass Spectrum AP^+ : Found 226 (MH^+). $\text{C}_{11}\text{H}_{15}\text{NSO}_2$ requires 225.

Description 31

10 *trans*-3-(2-(1-(4-(*N*-*tert*-Butyloxycarbonyl)amino)cyclohexyl)ethyl)-7-methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine

A solution of 7-methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine (1.0 g, 4.67 mmol) and *trans*-(1-(4-*N*-*tert*-butyloxycarbonyl)amino)cyclohexylacetaldehyde (0.8 g, 3.34 mmol) in dichloroethane (20 ml) was stirred at room temperature for 5 min before sodium triacetoxyborohydride (0.95 g, 4.49 mmol) was added in a single portion. After stirring at
15 room temperature for 48 h, the reaction mixture was partitioned between water (50 ml) and dichloromethane (100 ml). The aqueous layer was separate, re-extracted with dichloromethane (2 x 50 ml) and the combined organic layers dried (Na_2SO_4). The solvent was then removed *in vacuo* to give a pale yellow solid which was purified by column chromatography (silica gel; ethyl acetate : methanol; 9 : 1) to give the title
20 compound (1.35 g, 90 %) as a colourless solid.

^1H NMR (CDCl_3): 0.99 - 1.14 (4H, m), 1.23 - 1.29 (1H, m), 1.41 - 1.46 (2H, m), 1.46 (9H, s), 1.73 - 1.79 (2H, m), 2.00 - 2.06 (2H, m), 2.50 (2H, t, $J = 7.6$ Hz), 2.62 - 2.65 (4H, m), 2.99 - 3.02 (4H, m), 3.05 (3H, s), 3.38 (1H, br s), 4.38 (1H, br s), 7.27 - 7.30 (1H, d),
25 7.67 - 7.74 (2H, m).

Mass spectrum: AP^+ Found: 351 ($[\text{M}-\text{BOC}]\text{H}^+$). $\text{C}_{24}\text{H}_{38}\text{N}_2\text{SO}_4$ requires 450.

Description 32

30 *trans*-3-(2-(1-(4-Amino)cyclohexyl)ethyl)-7-methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine

A solution of *trans*-3-(2-(1-(4-*N*-*tert*-butyloxycarbonyl)amino)cyclohexyl)ethyl-7-methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine (1.3 g, 2.89 mmol) in dichloromethane (24 ml) and trifluoroacetic acid (6 ml) were stirred at room temperature
35 for 2 h. The reaction mixture was then concentrated *in vacuo* and the residue partitioned

between water (60 ml) and ethyl acetate (20 ml). The aqueous layer was separated, extracted with ethyl acetate (30 ml) and then basified to pH = 14 with 40 % sodium hydroxide. The oily suspension was then extracted with ethyl acetate (3 x 60 ml) and the combined organic layers dried (Na₂SO₄). The solvent was evaporated *in vacuo* to give the title compound (1.01 g, 100 %) as an off-white solid.

¹H NMR (CDCl₃) δ: 0.90 - 1.12 (4H, m), 1.15 - 1.22 (1H, m), 1.35 - 1.40 (2H, m), 1.72 - 1.78 (2H, m), 1.82 - 1.90 (2H, m), 2.45 - 2.52 (2H, m), 2.55 - 2.62 (5H, m), 2.98 - 3.02 (4H, m), 3.04 (3H, s), 7.27 (1H, d, J = 7.8 Hz), 7.56 (1H, s), 7.68 (1H, d).

Mass spectrum: AP⁺ 351 (MH⁺): C₁₉H₃₀N₂SO₂ requires 350.

Description 33

3-(*tert*-Butyloxycarbonyl)-7-(5-(3-methyl)isoxazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine

A mixture of 7-acetyl-3-(*tert*-butyloxycarbonyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine (6.18 g, 21.4 mmol) and dimethylacetamide dimethylacetal (8 ml) was stirred at 125 °C. Methanol by-product was removed by means of a Dean-Stark apparatus. After 8 h, excess dimethyl acetamide dimethyl acetal was evaporated *in vacuo* to give a thick oily residue. Absolute ethanol (20 ml) and hydroxylamine hydrochloride (2.53 g, 36.4 mmol) were added and the resulting mixture was heated under reflux for 2 h. The ethanol was removed *in vacuo* and the crude product residue was purified by silica gel chromatography eluting with 10 - 100 % ethyl acetate in hexane to give the title compound as a colourless oil (6.1 g, 87 %).

Mass spectrum (API⁺): Found 351 (MNa⁺). C₁₉H₂₄N₂O₃ requires 328.

¹H NMR (CDCl₃) δ: 1.49 (9H, s), 2.35 (3H, s), 2.90 - 3.00 (4H, m), 3.50 - 3.65 (4H, m), 6.31 (1H, s), 7.21 (1H, d, J = 8 Hz), 7.50 - 7.53 (2H, m).

Description 34

7-(5-(3-Methyl)isoxazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine

A solution of 3-(*tert*-butyloxycarbonyl)-7-(5-(3-methyl)isoxazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine (5.1 g, 15.6 mmol) in CH₂Cl₂ (30 ml) and trifluoroacetic acid (10 ml) was heated under reflux for 2 h. Solvent was evaporated *in vacuo* and the residue partitioned between diethyl ether (150 ml) and water (150 ml). The aqueous phase was

saturated with potassium carbonate then extracted with CH₂Cl₂ (2 x 200 ml). The combined organic extracts were dried and evaporated *in vacuo* to give the title compound (3.15 g, 88 %).

- 5 Mass spectrum (API⁺): Found 229 (MH⁺). C₁₄H₁₆N₂O requires 228.

¹H NMR (CDCl₃) δ: 1.80 (1H, br s), 2.34 (3H, s), 2.90 - 3.10 (8H, m), 6.30 (1H, s), 7.17 (1H, d, J = 8 Hz), 7.40 - 7.55 (2H, m).

10 **Description 35**

***trans*-3-(2-(1-(4-*N*-*tert*-Butyloxycarbonyl)amino)cyclohexyl)ethyl-7-(5-(3-methyl)isoxazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

Prepared from 7-(5-(3-methyl)isoxazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine in a manner similar to Description 2, in 92% yield.

15

Mass spectrum (API⁺): Found 454 (MH⁺). C₂₇H₃₉N₃O₃ requires 453.

¹H NMR (CDCl₃) δ: 1.00 - 1.10 (4H, m), 1.15 - 1.25 (1H, m), 1.44 (9H, s), 1.55 - 1.70 (2H, m), 1.70 - 1.85 (2H, m), 1.95 - 2.05 (2H, m), 2.34 (3H, s), 2.45 - 2.55 (2H, m), 2.55 - 2.70 (4H, m), 2.90 - 3.00 (4H, m), 3.35 (1H, m), 4.30 - 4.40 (1H, m), 6.30 (1H, s), 7.16 (1H, d, J = 8 Hz), 7.45 - 7.55 (2H, m).

20

Description 36

***trans*-3-(2-(1-(4-Amino)cyclohexyl)ethyl-7-(5-(3-methyl)isoxazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

25

Prepared from *trans*-3-(2-(1-(4-*N*-*tert*-butyloxycarbonyl)amino)cyclohexyl)ethyl-7-(5-(3-methyl)isoxazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine in a manner similar to Description 3 in 99 % yield.

¹H NMR (CDCl₃) δ: 0.90 - 1.10 (4H, m), 1.15 - 1.25 (1H, m), 1.35 - 1.50 (4H, m), 1.70 - 1.80 (2H, m), 1.80 - 1.90 (2H, m), 2.34 (3H, s), 2.42 - 2.52 (2H, m), 2.55 - 2.72 (5H, m), 2.90 - 3.00 (4H, m), 6.30 (1H, s), 7.16 (1H, d, J = 8 Hz), 7.45 - 7.55 (2H, m).

30

Description 37**3-(*tert*-Butyloxycarbonyl)-7-methanesulphonyloxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

5

A solution of 3-(*tert*-butyloxycarbonyl)-7-hydroxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine (3.0 g, 0.011 mol), methanesulphonylchloride (1.44 g, 0.013 mol), triethylamine (1.27 g, 0.013 mol) and dichloromethane (50 ml) was stirred at room temperature for 18 h. The reaction mixture was then partitioned between dichloromethane (50 ml) and a saturated solution of sodium hydrogen carbonate (50 ml). The organic layer was separated, washed with water (50 ml) and then dried (Na₂SO₄). The solvent was then evaporated *in vacuo* to give the title compound (3.85 g, 99 %) as a pale yellow oil.

¹H NMR (CDCl₃) δ : 1.48 (9H, s), 2.86 - 2.92 (4H, m), 3.13 (3H, s), 3.53 - 3.56 (4H, m), 7.00 - 7.03 (2H, m), 7.13 - 7.16 (1H, m).

Mass spectrum (AP⁺) : Found 242 [M-BOC]H⁺. C₁₆H₂₃NSO₅ requires 341.

20 Description 38**7-Methanesulphonyloxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

A solution of 3-(*tert*-butyloxycarbonyl)-7-methanesulphonyloxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine (3.8 g, 0.011 mol), trifluoroacetic acid (3.76 g, 0.033 mol) and dichloromethane (50 ml) was heated at 50 °C for 5h. The solvents were then evaporated *in vacuo* and the residue partitioned between water (200 ml) and ethyl acetate (150 ml). The aqueous layer was removed and washed with ethyl acetate (100 ml) and then basified to pH 14 with 40% sodium hydroxide. The suspension was then extracted with ethyl acetate (3 x 150 ml) and the combined organic layers dried (Na₂SO₄). The solvents were evaporated *in vacuo* to give the title compound (2.15 g, 80 %) as a colourless oil.

¹H NMR (CDCl₃) δ : 2.88 - 3.00 (8H, m), 3.13 (3H, s), 6.99 - 7.03 (2H, m), 7.12 (1H, d).

Mass spectrum (AP⁺) : Found 242 (MH)⁺. C₁₁H₁₅NSO₃ requires 241.

Description 39**40 *trans*-3-(2-(1-(4-*N*-*tert*-Butyloxycarbonyl)amino)cyclohexyl)ethyl-7-methanesulphonyloxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

A mixture of 7-(methanesulphonyloxy)-2,3,4,5-tetrahydro-1*H*-3-benzazepine (2.0 g, 8.3 mmol) and *trans*-2-(1-(4-*N*-*tert*-butyloxycarbonyl)amino)cyclohexyl acetaldehyde (1.37

g, 5.7 mmol) in dichloroethane (30 ml) was stirred at room temperature for 5 min, before sodium triacetoxymethylborohydride (1.69 g, 7.98 mmol) was added in a single portion. After stirring at room temperature for 48 h, a saturated solution of sodium hydrogen carbonate (50 ml) was added and the two layers separated. The aqueous layer was extracted with dichloromethane (3 x 60 ml) and the combined organic layers were dried (Na₂SO₄). The solvent was then evaporated *in vacuo* and the residue purified by column chromatography (silica gel, ethyl acetate) to give the title compound (2.54 g, 95 %) as a white solid.

¹H NMR (CDCl₃) δ : 0.9 - 1.25 (7H, m), 1.44 (9H, s), 1.70 - 1.80 (2H, m), 1.90 - 2.05 (2H, m), 2.42 - 2.50 (2H, m), 2.55 - 2.65 (4H, m), 2.88 - 2.95 (4H, m), 3.12 (3H, s), 3.36 (1H, br s), 4.34 (1H, br s), 6.98 - 7.02 (2H, m), 7.08 - 7.12 (1H, d).

Mass spectrum (AP⁺) : Found 467 [MH⁺]. C₂₄H₃₈N₂SO₅ requires 466.

15

Description 40

trans-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-methanesulphonyloxy-2,3,4,5-tetrahydro-1H-3-benzazepine

20

A solution of *trans*-3-(2-(1-(4-*N*-*tert*-butyloxycarbonyl)amino)cyclohexyl)ethyl-7-methanesulphonyloxy-2,3,4,5-tetrahydro-1H-3-benzazepine, trifluoroacetic acid (8 ml) and dichloromethane (32 ml), were stirred at room temperature for 2 h, under argon. The solvents were then evaporated *in vacuo* and the residue partitioned between water (150 ml) and ethyl acetate (60 ml). The aqueous layer was removed and washed with ethyl acetate (50 ml). The aqueous layer was then basified to pH 14 with 40% sodium hydroxide. The suspension was then extracted with ethyl acetate (3 x 80 ml) and the combined organic layers dried (Na₂SO₄). The solvents were evaporated *in vacuo* to give the title compound (1.78 g, 93 %) as an oil which crystallised on standing.

30

¹H NMR (CDCl₃) δ : 0.95 - 1.45 (7H, m), 1.70 - 1.80 (2H, m), 1.80 - 1.90 (2H, m), 2.49 (2H, t, J = 7.8 Hz), 2.55 - 2.65 (5 H, m), 2.88 - 2.95 (4H, m), 3.12 (3H, s), 6.99 - 7.02 (2H, m), 7.11 (1H, d, J = 8 Hz).

35


Mass Spectrum (AP⁺): Found 367 (MH⁺). C₁₉H₃₀N₂SO₃ requires 366.

Examples

The Compounds of Examples tabulated below (Tables 1 – 3) were all prepared using the following general method:-

- 5 A mixture of the appropriate *trans*-2-(2-(1-(4-amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine (0.35 mmol), the appropriate acid (0.35 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (0.35 mmol), 1-hydroxybenzotriazole (catalytic amount) and dichloromethane (5ml) was shaken for 16h.
- 10 Saturated sodium bicarbonate (4ml) was then added and the mixture shaken for 0.25h. Chromatography of the organic layer on silica with 50 - 100% ethyl acetate in hexane and 0 - 10% methanol in ethyl acetate gradient elution gave the title compounds.

Table 1.



15

Example	R ¹	A	Characterising Data
			Mass Spectrum (API ⁺); ¹ H NMR (CDCl ₃)
1	H	4-Quinolinylnyl	Found: 428 (MH ⁺). C ₂₈ H ₃₃ N ₃ O requires 427. δ: 1.06 – 1.37 (5H, m), 1.47 (2H, m), 1.78 (2H, m), 2.19 (2H, m), 2.51 (2H, m), 2.64 (4H, m), 2.93 (4H, m), 4.03 (1H, m), 5.90 (1H, d, J = 8 Hz), 7.10 (4H, m), 7.40 (1H, d, J = 4 Hz), 7.60 (1H, m), 7.75 (1H, m), 8.15 (2H, m), 8.91 (1H, d, J = 4 Hz).
2	H	<i>trans</i> -CH=CHC ₆ H ₄ (3-SO ₂ Me)	Found: 481 (MH ⁺). C ₂₈ H ₃₆ N ₂ O ₃ S requires 480. δ: 1.01 – 1.33 (5H, m), 1.45 (2H, m), 1.81 (2H, m), 2.02 (2H, m), 2.50 (2H, m), 2.63 (4H, m), 2.93 (4H, m), 3.07 (3H, s), 3.84 (1H, m), 5.64 (1H, d, J = 8 Hz), 6.49 (1H, d, J = 16 Hz), 7.09 (4H, m), 7.64 (3H, m), 7.89 (1H, d, J = 8 Hz), 8.09 (1H, s).
3	H	<i>trans</i> -	Found: 421 (MH ⁺). C ₂₇ H ₃₃ FN ₂ O

		$\text{CH=CHC}_6\text{H}_4(4\text{-F})$	requires 420. δ : 1.00 – 1.33 (5H, m), 1.43 (2H, m), 1.79 (2H, m), 2.04 (2H, m), 2.49 (2H, m), 2.62 (4H, m), 2.93 (4H, m), 3.86 (1H, m), 5.51 (1H, d, J = 8 Hz), 6.26 (1H, d, J = 16 Hz), 7.11 (6H, m), 7.47 (2H, m), 7.56 (1H, d, J = 16 Hz).
4	H	2-Indolyl	Found: 416 (MH ⁺). C ₂₇ H ₃₃ N ₃ O requires 415. δ : 1.05 – 1.37 (5H, m), 1.46 (2H, m), 1.82 (2H, m), 2.12 (2H, m), 2.52 (2H, m), 2.63 (4H, m), 2.92 (4H, m), 3.96 (1H, m), 5.98 (1H, d, J = 8 Hz), 6.79 (1H, m), 7.14 (5H, m), 7.26 (1H, m), 7.41 (1H, m), 7.64 (1H, d, J = 8 Hz), 9.35 (1H, br s).
5	H	-C ₆ H ₄ (3-(3-pyridyl))	Found: 454 (MH ⁺). C ₃₀ H ₃₅ N ₃ O requires 453. δ : 1.08 – 1.37 (5H, m), 1.46 (2H, m), 1.84 (2H, m), 2.12 (2H, m), 2.51 (2H, m), 2.63 (4H, m), 2.92 (4H, m), 3.95 (1H, m), 6.02 (1H, d, J = 8 Hz), 7.11 (4H, m), 7.38 (1H, m), 7.54 (1H, t, J = 8 Hz), 7.72 (2H, m), 7.91 (1H, m), 7.98 (1H, s), 8.62 (1H, m), 8.86 (1H, d, J = 2 Hz).
6	H	-CH ₂ Ph	Found 391 (MH ⁺). C ₂₆ H ₃₄ N ₂ O requires 390. δ : 0.87 – 1.14 (5H, m), 1.40 (2H, m), 1.68 (2H, m), 1.89 (2H, m), 2.45 (2H, m), 2.49 (4H, m), 2.89 (4H, m), 3.54 (2H, s), 3.68 (1H, m), 5.14 (1H, m), 7.10 (4H, m), 7.30 (5H, m).
7	H	-CH ₂ (3-indolyl)	Found: 430 (MH ⁺). C ₂₈ H ₃₅ N ₃ O requires 429. δ : 0.77 – 1.13 (5H, m), 1.36 (2H, m), 1.64 (2H, m), 1.84 (2H, m), 2.42 (2H, m), 2.57 (4H, m), 2.98 (4H, m), 3.70 (3H, m), 5.46

			(1H, d, J = 8 Hz), 7.12 (7H, m), 7.39 (1H, d, J = 8 Hz), 7.53 (1H, d, J = 8 Hz), 8.24 (1H, br s).
8	H	-CH ₂ (4-quinolyl)	<p>Found: 442 (MH⁺) C₂₉H₃₅N₃O requires 441.</p> <p>δ: 0.80 – 1.15 (5H, m), 1.35 – 1.45 (2H, m), 1.65 – 1.75 (2H, m), 1.80 – 1.90 (2H, m), 2.45 – 2.52 (2H, m), 2.60 – 2.70 (4H, m), 2.85 – 3.00 (4H, m), 3.70 (1H, m), 3.99 (2H, s), 5.13 (1H, d, J = 8 Hz), 7.00 – 7.16 (4H, m), 7.33 (1H, d, J = 3 Hz), 7.60 (1H, m), 7.75 (1H, m), 7.98 (1H, d, J = 8 Hz), 8.14 (1H, d, J = 8 Hz), 8.88 (1H, d, J = 3 Hz).</p>
9	6-OMe	<i>trans</i> -CH=CHC ₆ H ₄ (4-F)	<p>Found: 451 (MH⁺), C₂₈H₃₅FN₂O₂ requires 450.</p> <p>δ: 1.10 – 1.30 (5H, m), 1.40 – 1.50 (2H, m), 1.75 – 1.85 (2H, m), 2.00 – 2.10 (2H, m), 2.40 – 2.52 (2H, m), 2.55 – 2.70 (4H, m), 2.85 – 3.10 (4H, m), 3.80 (3H, s), 3.86 (1H, m), 5.37 (1H, d, J = 8 Hz), 6.26 (1H, d, J = 15 Hz), 6.74 (2H, m), 7.06 (3H, m), 7.47 (2H, dd, J = 9 Hz, 6 Hz), 7.54 (1H, d, J = 15 Hz).</p>
10	6-OMe	4-Quinolyl	<p>Found: 458 (MH⁺), C₂₉H₃₅N₃O₂ requires 457</p> <p>δ: 1.05 – 1.35 (5H, m), 1.55 – 1.69 (2H, m), 1.70 – 1.80 (2H, m), 2.10 – 2.24 (2H, m), 2.90 – 3.35 (10H, m), 3.80 (3H, s), 4.00 (1H, m), 5.85 (1H, d, J = 8 Hz), 6.75 (2H, m), 7.13 (1H, t, J = 8 Hz), 7.41 (1H, d, J = 4 Hz), 7.60 (1H, m), 7.78 (1H, m), 8.16 (2H, m), 8.94 (1H, d, J = 4 Hz).</p>
11	6-OMe	3-(pyrrolo[2,3-b]pyridyl)	<p>Found: 447 (MH⁺), C₂₇H₃₄N₄O₂ requires 446.</p> <p>δ: 1.30 – 1.35 (5H, m), 1.45 – 1.55 (2H, m), 1.75 – 1.80 (2H, m), 2.10 – 2.20 (2H,</p>

			m), 2.55 – 2.85 (6H, m), 2.90 – 3.20 (4H, m), 3.80 (3H, s), 3.96 (1H, m), 5.65 (1H, d, J = 8 Hz), 6.74 (2H, m), 7.10 (1H, t, J = 8 Hz), 7.20 (1H, m), 7.80 (1H, s), 8.35 (2H, m), 9.45 (1H, br s).
30	7-CN	-CH ₂ Ph(2,5-diF)	Found: 452 (MH ⁺); C ₂₇ H ₃₁ F ₂ N ₃ O requires 451. δ: 0.80 - 1.00 (4H, m), 1.10 (1H, m), 1.40 - 1.50 (2H, m), 1.55 - 1.65 (2H, m), 1.80 - 1.90 (2H, m), 2.70 - 2.80 (2H, m), 2.90 - 3.20 (8H, m), 3.48 (2H, s), 3.60 (1H, m), 5.28 (1H, d, J = 8 Hz), 6.90 - 7.05 (3H, m), 7.21 (1H, d, J = 8 Hz), 7.40 (1H, s), 7.45 (1H, d, J = 6 Hz).
31	7-CN	-CH ₂ (2-naphthyl)	Found: 466 (MH ⁺); C ₃₁ H ₃₅ N ₃ O requires 465. δ: 0.80 - 1.20 (5H, m), 1.30 - 1.40 (2H, m), 1.65 - 1.75 (2H, m), 1.85 - 1.90 (2H, m), 2.35 - 2.50 (2H, m), 2.50 - 2.60 (4H, m), 2.85 - 3.00 (4H, m), 3.60 - 3.85 (3H, m), 5.16 (1H, d, J = 9 Hz), 7.10 (1H, d, J = 9 Hz), 7.30 - 7.40 (3H, m), 7.40 - 7.60 (2H, m), 7.70 (1H, s), 7.80 - 7.90 (3H, m).
32	7-CN	<i>trans</i> -CH=CHC ₆ H ₃ (2,4-diF)	Found: 464 (MH ⁺); C ₂₈ H ₃₁ N ₃ OF ₂ requires 463. δ: 1.02 - 1.30 (5H, m), 1.40 - 1.48 (2H, m), 1.78 - 1.82 (2H, m), 2.03 - 2.06 (2H, m), 2.50 (2H, t, J = 8 Hz), 2.62 - 2.66 (4H, m), 2.93 - 3.00 (4H, m), 3.79 - 3.91 (1H, m), 5.72 (1H, d, J = 8 Hz), 6.43 (1H, d, J = 16 Hz), 6.75 - 6.95 (2H, m), 7.18 (1H, d, J = 8 Hz), 7.38 - 7.47 (3H, m), 7.59 (1H, d, J = 16 Hz).
33	7-CN	<i>trans</i> -CH=CHC ₆ H ₃ (2,5-diF)	Found: 464 (MH ⁺); C ₂₈ H ₃₁ N ₃ OF ₂ requires 463. (CD ₃ OD) δ: 1.06 - 1.40 (5H, m), 1.60 - 1.74 (2H, m), 1.80 - 1.85 (2H, m), 1.92 -

			1.97 (2H, m), 2.90 - 3.40 (8H, m), 3.64 - 3.79 (3H, m), 6.64 (1H, d, J = 16 Hz), 7.09 - 7.18 (2H, m), 7.29 - 7.39 (2H, m), 7.52 - 7.58 (3H, m).
34	7-CN	<i>trans</i> - CH=CHC ₆ H ₄ (3-F)	Found: 446 (MH ⁺); C ₂₈ H ₃₂ N ₃ O requires 445. δ: 1.03 - 1.35 (5H, m), 1.40 - 1.48 (2H, m), 1.75 - 2.10 (4H, m), 2.51 (2H, t, J = 8 Hz), 2.61 - 2.65 (4H, m), 2.93 - 2.99 (4H, m), 3.72 - 3.88 (1H, m), 6.36 (1H, d, J = 18 Hz), 7.04 (1H, m), 7.17 - 7.45 (7H, m), 7.55 (1H, d, J = 16 Hz)
35		-CH ₂ CH ₂ C ₆ H ₅	Found: 430 (MH ⁺); C ₂₈ H ₃₅ N ₃ O requires 429. δ: 0.90 - 1.25 (5H, m), 1.30 - 1.45 (2H, m), 1.65 - 1.95 (4H, m), 2.35 - 2.50 (4H, m), 2.55 - 2.65 (4H, m), 2.90 - 3.04 (6H, m), 3.66 - 3.69 (1H, m), 5.09 (1H, d, J = 8 Hz), 7.15 - 7.35 (6H, m), 7.37 - 7.42 (2H, m).
36	7-CN	<i>trans</i> - CH=CHC ₆ H ₄ (2-F)	Found: 446 (MH ⁺), C ₂₈ H ₃₂ FN ₃ O requires 445. δ: 1.00 - 1.30 (5H, m), 1.40 - 1.50 (2H, m), 1.75 - 1.85 (2H, m), 2.00 - 2.10 (2H, m), 2.45 - 2.50 (2H, m), 2.60 - 2.70 (4H, m), 2.85 - 3.00 (4H, m), 3.80 - 3.95 (1H, m), 5.43 (1H, d, J = 8 Hz), 6.47 (1H, d, J = 16 Hz), 7.00 - 7.20 (3H, m), 7.26 - 7.35 (1H, m), 7.30 - 7.55 (3H, m), 7.66 (1H, d, J = 16 Hz).
37	7-CN	8-(1,4-dihydro-4-oxo)-quinolinyl	Found: 469 (MH ⁺); C ₂₉ H ₃₂ N ₄ O ₂ requires 468. δ: 1.05 - 1.35 (5H, m), 1.40 - 1.50 (2H, m), 1.80 - 1.90 (2H, m), 2.10 - 2.20 (2H, m), 2.45 - 2.55 (2H, m), 2.60 - 2.70 (4H, m), 2.90 - 3.00 (4H, m), 3.95 (1H, m), 6.30 - 6.40 (2H, m), 7.18 (1H, d, J = 8 Hz), 7.32 (1H, t, J = 8 Hz), 7.37 (1H, s), 7.41 (1H, d,

			J = 8 Hz), 7.67 (1H, t, J = 5 Hz), 7.80 (1H, d, J = 7 Hz), 8.52 (1H, d, J = 8 Hz), 12.50 (1H, br s).
38	7-CN	2-naphthyl	Found: 452 (MH ⁺); C ₃₀ H ₃₃ N ₃ O requires 451. δ: 1.09 - 1.26 (5H, m), 1.42 - 1.50 (2H, m), 1.80 - 1.87 (2H, m), 2.10 - 2.25 (2H, m), 2.40 - 2.54 (2H, m), 2.61 - 2.70 (4H, m), 2.92 - 2.99 (4H, m), 3.95 - 4.00 (1H, m), 5.81 (1H, d, J = 8 Hz), 7.18 (1h, d, J = 8 Hz), 7.30 - 7.59 (6H, m), 7.84 - 7.95 (2H, m), 8.20 - 8.29 (1H, m).
39	7-CN	<i>trans</i> -CH=CHC ₆ H ₄ (2-OMe)	Found: 458 (MH ⁺); C ₂₉ H ₃₅ N ₃ O ₂ requires: 457. (DMSO-d ₆) δ: 0.99 - 1.07 (2H, m), 1.15 - 1.28 (3H, m), 1.35 - 1.40 (2H, m), 1.78 - 1.88 (4H, m), 2.46 (2H, t, J = 7 Hz), 2.58 (4H, m, obscured by DMSO), 2.90 - 2.96 (4H, m), 3.58 - 2.64 (1H, m), 3.87 (3H, s), 6.64 (1H, d, J = 16 Hz), 6.99 (1H, t, J = 7 Hz), 7.09 (1H, d, J = 8 Hz), 7.34 - 7.39 (2H, m), 7.49 - 7.51 (1H, m), 7.58 - 7.66 (3H, m), 7.93 - 7.96 (1H, m).
40	7-CN	<i>trans</i> -CH=CHC ₆ H ₄ (3-OMe)	Found: 458 (MH ⁺); C ₂₉ H ₃₅ N ₃ O ₂ requires 457. (DMSO-d ₆) δ: 0.80 - 0.98 (2H, m), 1.00 - 1.20 (3H, m), 1.20 - 1.35 (2H, m), 1.78 - 1.82 (2H, m), 1.88 - 1.92 (2H, m), 2.49 (2H, t, J = 8 Hz), 2.51 - 2.60 (4H, m, obscured by DMSO), 2.90 - 3.00 (4H, m), 3.62 (1H, m), 3.83 (3H, s), 6.65 (1H, d, J = 16 Hz), 7.00 (1H, m), 7.15 - 7.17 (2H, m), 7.35 - 7.43 (3H, m), 7.61 - 7.63 (2H, m), 7.99 (1H, d, J = 8 Hz).
41	7-CN	<i>trans</i> -CH=CHC ₆ H ₄ (4-OMe)	Found: 458 (MH ⁺); C ₂₉ H ₃₅ N ₃ O ₂ requires: 457. (DMSO-d ₆) δ: 1.00 - 1.10 (2H, m), 1.15 -

			1.27 (3H, m), 1.38 - 1.41 (2H, m), 1.77 - 1.80 (2H, m), 1.85 - 1.87 (2H, m), 2.47 (2H, t, J = 7 Hz), 2.53 - 2.57 (4H, m, obscured by DMSO), 2.91 - 2.97 (4H, m), 3.58 - 3.65 (1H, m), 3.81 (3H, s), 6.48 (1H, d, J = 16 Hz), 7.00 (2H, m), 7.35 - 7.40 (2H, m), 7.51 (2H, m), 7.58 - 7.52 (2H, m), 7.88 (1H, d, J = 8 Hz).
42	7-CN	<i>trans</i> - CH=CHC ₆ H ₄ (2-COMe)	Found: 470 (MH ⁺); C ₃₀ H ₃₅ N ₃ O ₂ requires 469. (DMSO-d ₆) δ: 0.90 - 1.40 (7H, m), 1.60 - 2.90 (4H, m), 2.46 (2H, t, J = 8 Hz), 2.46 - 2.54 (4H, m, obscured by DMSO), 2.52 (3H, s), 2.80 - 3.00 (4H, m), 3.60 (1H, m), 6.47 (1H, d, J = 16 Hz), 7.35 (2H, d, J = 8 Hz), 7.40 - 7.63 (4H, m), 7.75 (1H, d, J = 16 Hz), 7.89 (1H, d, J = 8 Hz), 8.04 (1H, d, J = 8 Hz).
43	7-CN	<i>trans</i> - CH=CHC ₆ H ₄ (4-COMe)	Found: 470 (MH ⁺); C ₃₀ H ₃₅ N ₃ O ₂ requires: 469. (DMSO-d ₆): 0.90 - 1.40 (7H, m), 1.73 - 1.88 (4H, m), 2.43 (2H, t, J = 8 Hz), 2.47 - 2.54 (4H, m, obscured by DMSO), 2.59 (3H, s), 2.80 - 2.93 (4H, m), 3.50 - 3.70 (1H, m), 6.73 (1H, d, J = 16 Hz), 7.32 (1H, d, J = 7 Hz), 7.45 (1H, d, J = 16 Hz), 7.55 - 7.59 (2H, m), 7.68 (2H, d, J = 8 Hz), 7.98 (2H, m), 8.08 (1H, d, J = 8 Hz).
44	7-CN	<i>trans</i> - CH=CHC ₆ H ₄ (2-CN)	Found: 453 (MH ⁺); C ₂₉ H ₃₂ N ₄ O requires 452. (DMSO-d ₆) δ: 1.02 - 1.09 (2H, m), 1.10 - 1.35 (3H, m), 1.36 - 1.42 (2H, m), 1.78 - 1.81 (2H, m), 1.88 - 1.90 (2H, m), 2.47 (2H, t, J = 7 Hz), 2.50 - 2.59 (4H, m, obscured by DMSO), 2.92 - 2.97 (4H, m), 3.62 - 3.58 (1H, m), 6.85 (1H, d, J = 16 Hz), 7.36 (1H, d, J = 8 Hz), 7.58 - 7.67 (4H, m), 7.79 - 7.89 (2H, m), 7.92 - 7.95

			(1H, m), 8.22 - 8.25 (1H, m).
45	7-CN	<i>trans</i> - CH=CHC ₆ H ₄ (3-CN)	Found: 453 (MH ⁺); C ₂₉ H ₃₂ N ₄ O requires: 452. (DMSO-d ₆) δ: 0.94 - 1.38 (7H, m), 1.70 - 1.87 (4H, m), 2.43 (2H, t, J = 7 Hz), 2.46 - 2.59 (4H, m, obscured by DMSO), 2.85 - 2.97 (4H, m), 3.52 - 3.65 (1H, m), 6.72 (1H, d, J = 16 Hz), 7.32 (1H, d, J = 8 Hz), 7.42 (1H, d, J = 16 Hz), 7.55 - 7.62 (3H, m), 7.80 - 7.91 (2H, m), 8.02 (1H, s), 8.09 (1H, d, J = 8 Hz).
46	7-CN	-C ₆ H ₄ (3-(5-(3-methyl)isoxazolyl))	Found: 483 (MH ⁺); C ₃₀ H ₃₄ N ₄ O ₂ requires 482. (DMSO-d ₆) δ: 0.96 - 1.10 (2H, m), 1.23 - 1.50 (5H, m), 1.70 - 1.89 (4H, m), 2.31 (3H, s), 2.42 - 2.55 (6H, m, obscured by DMSO), 2.80 - 2.95 (4H, m), 3.75 (1H, m), 6.96 (1H, s), 7.33 (1H, d, J = 8 Hz), 7.50 - 7.60 (3H, m), 7.90 - 8.00 (2H, m), 8.28 (1H, m), 8.41 (1H, d, J = 8 Hz).
47	7-CN	7-(1,2-dihydro-2-oxo)quinolinyl	Found: 469 (MH ⁺); C ₂₉ H ₃₂ N ₄ O ₂ requires 468. (DMSO-TFA) δ: 0.98 - 1.45 (5H, m), 1.56 - 1.63 (1.56 - 1.63 (2H, m), 1.75 - 1.89 (4H, m), 2.95 - 3.32 (8H, m), 3.65 - 3.85 (3H, m), 5.67 (1H, s), 6.60 (1H, d, J = 10 Hz), 7.41 (1H, d, J = 8 Hz), 7.59 - 7.70 (4H, m), 7.75 (1H, s), 7.90 (1H, d, J = 10 Hz), 8.32 (1H, d), 9.69 (1H, s).
48	7-CN	<i>cis</i> -CH=CHC ₆ H ₅	Found: 428 (MH ⁺); C ₂₈ H ₃₃ N ₃ O requires 427. δ: 0.80 - 1.15 (5H, m), 1.30 - 1.40 (2H, m), 1.65 - 1.75 (2H, m), 1.80 - 1.95 (2H, m), 2.40 - 2.50 (2H, m), 2.55 - 2.65 (4H, m), 2.85 - 3.00 (4H, m), 3.75 (1H, m), 5.25 (1H, d, J = 8 Hz), 5.98 (1H, d, J = 12.5 Hz), 6.76 (1H, d, J = 12.5 Hz), 7.17 (1H, d, J =

			8 Hz), 7.30 - 7.45 (7H, m).
49	7-CN	<i>trans</i> -CH=CH(2-pyridyl)	<p>Found: 429 (MH⁺); C₂₇H₃₂N₄O requires 428.</p> <p>δ (DMSO+TFA): 0.90 - 1.30 (5H, m), 1.55 - 1.70 (2H, m), 1.70 - 1.80 (2H, m), 1.80 - 1.90 (2H, m), 2.90 - 3.30 (8H, m), 3.50 - 3.80 (3H, m), 7.08 (1H, d, J = 16 Hz), 7.40 - 7.50 (2H, m), 7.55 - 7.60 (1H, m), 7.65 - 7.80 (3H, m), 8.05 (1H, m), 8.25 (1H, d, J = 8 Hz), 8.70 (1H, m), 9.70 (1H, br s).</p>
50	7-CN	<i>trans</i> -CH=CH(1-(4-fluoro)naphthyl)	<p>Found: 496 (MH⁺); C₃₂H₃₄FN₃O requires 495.</p> <p>δ: (DMSO-d₆ + TFA); 0.97 - 1.41 (5H, m), 1.63 (2H, m), 1.79 (2H, m), 1.90 (2H, m), 3.06 (2H, m), 3.23 (6H, m), 3.70 (3H, m), 6.64 (1H, d, J = 16 Hz), 7.47 (2H, m), 7.73 (5H, m), 8.12 (3H, m), 8.24 (1H, m).</p>
51	7-CN	<i>trans</i> -CH=CH(6-benzodioxanyl)	<p>Found: 486 (MH⁺); C₃₀H₃₅N₃O requires 485.</p> <p>δ: (DMSO d₆ + TFA): 0.93 - 1.33 (5H, m), 1.60 (2H, m), 1.81 (4H, m), 3.04 (2H, m), 3.17 (6H, m), 3.67 (3H, m), 4.26 (4H, s), 6.42 (1H, d, J = 16 Hz), 6.87 (1H, d, J = 9 Hz), 7.03 (2H, m), 7.27 (1H, d, J = 16 Hz), 7.46 (1H, d, J = 8 Hz), 7.73 (2H, m), 7.90 (1H, d, J = 8 Hz), 9.78 (1H, br s).</p>
52	7-CN	<i>trans</i> -CH=CH(3-indolyl[5-F])	<p>Found: 485 (MNa⁺); C₃₀H₃₃FN₄O requires 484.</p> <p>(DMSO-d₆) δ: 0.98 - 1.08 (2H, m), 1.11 - 1.28 (3H, m), 1.35 - 1.42 (2H, m), 1.75 - 1.80 (2H, m), 1.87 - 1.91 (2H, m), 2.47 (2H, t, J = 7 Hz), 2.52 - 2.59 (4H, m, obscured by DMSO), 2.89 - 2.94 (4H, m), 3.55 - 3.62 (1H, m), 6.56 (1H, d, J = 16 Hz), 7.03 - 7.09 (1H, m), 7.34 (1H, d, J = 8 Hz), 7.44 - 7.49 (1H, m), 7.56 - 7.75 (5H,</p>

			m), 7.80 (1H, s), 11.63 (1H, s).
53	7-CN	<i>trans</i> -CH=CH(6-benzimidazolyl[1-methyl])	Found: 482 (MH ⁺); C ₃₀ H ₃₅ N ₅ O requires 481. (DMSO-d ₆) δ: 0.98 - 1.07 (2H, m), 1.16 - 1.27 (3H, m), 1.30 - 1.40 (2H, m), 1.75 - 1.79 (2H, m), 1.84 - 1.89 (2H, m), 2.46 (2H, t, J = 7 Hz), 2.50 - 2.55 (4H, m, obscured by DMSO), 2.90 - 2.97 (4H, m), 3.60 - 3.66 (1H, m), 3.87 (3H, s), 6.64 (1H, d, J = 16 HZ), 7.34 (1H, d, J = 8 Hz), 7.41 - 7.45 (1H, m), 7.53 - 7.45 (1H, m), 7.53 - 7.61 (3H, m), 7.66 (1H, d, J = 8 Hz), 7.77 (1H, s), 7.91 - 7.94 (1H, m), 8.24 (1H, s).
54	7-CN	<i>trans</i> -CH=CH(7-benzofuranyl)	Found: 468 (MH ⁺); C ₃₀ H ₃₃ N ₃ O ₂ requires 467. (DMSO/TFA) δ: 1.02 - 1.43 (5H, m), 1.65 - 1.75 (2H, m), 1.75 - 2.00 (4H, m), 3.08 - 3.35 (8H, m), 3.65 - 3.80 (3H, m), 7.09 - 7.15 (2H, m), 7.36 (1H, t); 7.48 - 7.56 (2H, m), 7.66 (1H, d, J = 15.8 Hz), 7.73 - 7.80 (3H, m), 8.15 (1H, d, J = 2.2 Hz), 8.25 (1H, d).
55	7-CN	<i>trans</i> -CH=CH(5-indolyl[3-methyl])	Found: 481 (MH ⁺) C ₃₁ H ₃₆ N ₄ O requires 480. (DMSO/TFA) δ: 0.95 - 1.35 (5H, m), 1.55 - 1.70 (2H, m), 1.70 - 1.95 (4H, m), 2.27 (3H, s), 2.95 - 3.30 (8H, m), 3.55 - 3.80 (3H, m), 6.52 (1H, d, J = 15.7 Hz), 7.14 (1H, s), 7.33 (2H, m), 7.46 (2H, m), 7.71 (3H, m), 7.84 (1H, d), 9.82 (1H, br s).
56	7-CN	<i>trans</i> -CH=CH(6-(2,3-dihydro-2-oxo)indolyl)	Found: 483 (MH ⁺); C ₃₀ H ₃₄ N ₄ O ₂ requires 482. (DMSO) δ: 0.87 - 1.36 (7H, m), 1.74 - 1.90 (4H, m), 2.44 (2H, t, J = 7.2 Hz), 2.50 - 2.65 (4H, m, under DMSO), 2.80 - 2.95 (4H, m), 3.49 (2H, s), 3.57 - 3.70 (1H, m), 6.54 (1H, d, J = 15.8 Hz), 6.95 (1H, s),

			7.09 (1H, d, J = 8), 7.22 (1H, d, J = 7.5), 7.27 - 7.39 (2H, m), 7.56 - 7.60 (2H, m), 7.92 (1H, d).
57	7-CN	-CH ₂ (2-benzofuranyl)	Found: 456 (MH ⁺); C ₂₉ H ₃₃ N ₃ O ₂ requires 455. (DMSO) δ: 0.95 - 1.23 (5H, m), 1.31 - 1.36 (2H, m), 1.72 - 1.82 (4H, m), 2.42 (2H, t, J = 7.4 Hz), 2.49 - 2.53 (4H, m, under DMSO), 2.88 - 2.93 (4H, m), 3.47 - 3.52 (1H, m), 3.63 (2H, s), 6.65 (1H, s), 7.19 - 7.24 (2H, m), 7.31 (1H, d, J = 7.7 Hz), 7.49 (1H, d, J = 7.9 Hz), 7.55 - 7.59 (3H, m), 8.02 (1H, d).
58	7-CN	<i>trans</i> -CH=CH(4-indolyl[2-methyl])	Found: 479 (MH ⁺); C ₃₁ H ₃₆ NO ₄ requires 480. (DMSO+TFA) δ: 0.83 - 1.25 (5H, m), 1.48 - 1.55 (2H, m), 1.63 - 1.67 (2H, m), 1.75 - 1.80 (2H, m), 2.31 (3H, s), 2.85 - 3.20 (8H, m), 3.45 - 3.65 (3H, m), 6.36 (1H, s), 6.62 (1H, d, J = 16 Hz), 6.89 (1H, t, J = 8 Hz), 7.00 (1H, d, J = 7 Hz), 7.18 (1H, d, J = 8 Hz), 7.34 (1H, d, J = 8 Hz), 7.49 (1H, d, J = 16 Hz), 7.55 - 7.62 (2H, m), 7.90 (1H, d, J = 8 Hz), 9.75 (1H, b s), 11.09 (1H, s).
59	7-CN	<i>trans</i> -CH=CH(5-benzimidazolyl)	Found: 466 (MH ⁺); C ₂₉ H ₃₃ N ₅ O requires 467. (DMSO+TFA) δ: 1.02 - 1.35 (5H, m), 1.57 - 1.61 (2H, m), 1.75 - 1.78 (2H, m), 1.90 - 1.93 (2H, m), 3.00 - 3.30 (8H, m), 3.65 - 3.70 (3H, m), 6.66 and 6.73 (1H, 2 x d, J = 16 Hz), 7.43 (1H, d, J = 8 Hz), 7.60 - 8.08 (6H, m), 8.00 and 8.06 (1H, 2 x s), 9.59 (1H, m), 9.88 (1H, b s).
60	7-CN	<i>trans</i> -CH=CHC ₆ H ₅	Found: 428 (MH ⁺); C ₂₈ H ₃₃ N ₃ O requires 427. (DMSO-d ₆ +TFA) δ: 0.96 - 1.36 (5H, m), 1.62 (2H, m), 1.81 (4H, m), 3.05 (2H, m),

			3.18 (6H, m), 3.67 (3H, m), 6.60 (1H, d, J = 16 Hz), 7.27 - 7.59 (7H, m), 7.72 (2H, m), 7.99 (1H, d, J = 8 Hz), 9.72 (1H, br s).
61	7-CN	<i>trans</i> - CH=CHC ₆ H ₃ (2,3-methylenedioxy)	Found: 472 (MH ⁺); C ₂₉ H ₃₃ N ₃ O ₃ requires 471. (DMSO-d ₆ +TFA) δ: 0.94 - 1.32 (5H, m), 1.61 (2H, m), 1.82 (4H, m), 3.03 (2H, m), 3.18 (6H, m), 3.64 (3H, m), 6.13 (2H, s), 6.71 (1H, d, J = 16 Hz), 6.94 (3H, m), 7.32 (1H, d, J = 16 Hz), 7.46 (1H, d, J = 8 Hz), 7.71 (2H, m), 8.09 (1H, d, J = 8 Hz), 9.75 (1H, br s).
62	7-CN	<i>trans</i> - CH=CHC ₆ H ₄ (3-(1-(2-oxo)pyrrolidinyl))	Found: 511 (MH ⁺); C ₃₂ H ₃₈ N ₄ O ₂ requires 510. (DMSO-d ₆) δ: 0.9 - 1.28 (5H, m), 1.35 (2H, m), 1.79 (4H, m), 2.07 (2H, m), 2.48 (8H, m), 2.91 (4H, m), 3.59 (1H, m), 3.86 (2H, t, J = 7 Hz), 6.60 (1H, d, J = 16 Hz), 7.34 (4H, m), 7.60 (3H, m), 7.89 (1H, m), 7.99 (1H, d, J = 8 Hz)...
63	7-CN	-CH ₂ (2-indolyl)	Found: 455 (MH ⁺); C ₂₉ H ₃₄ N ₄ O requires 454. (DMSO-d ₆) δ: 0.96 (2H, m), 1.15 (3H, m), 1.34 (2H, m), 1.76 (4H, m), 2.42 (2H, m), 2.50 (4H, m), 2.90 (4H, m), 3.45 (1H, m), 3.53 (2H, s), 6.17 (1H, m), 6.94 (2H, m), 7.32 (2H, m), 7.41 (1H, d, J = 8 Hz), 7.56 (2H, m), 7.87 (1H, d, J = 8 Hz), 10.85 (1H, br s).
64	7-CN	-CH ₂ (2-benzothiophenyl)	Found: 472 (MH ⁺); C ₂₉ H ₃₃ N ₃ SO requires 471. (DMSO) δ: 0.95 - 1.20 (5H, m), 1.31 - 1.35 (2H, m), 1.71 - 1.81 (4H, m), 2.42 (2H, t, J = 7.4 Hz), 2.50 - 2.53 (4H, m), 2.87 - 2.93 (4H, m), 3.44 - 3.48 (1H, m), 3.70 (2H, s), 7.19 (1H, s), 7.27 - 7.32 (3H, m), 7.55 - 7.58 (2H, m), 7.75 (1H, d, J = 7.4 Hz), 7.88

			(1H, d, J = 7.8 Hz), 8.04 (1H, m).
65		<i>trans</i> -CH=CH(2-thiophenyl[3-Br])	Found: 512 & 514 (MH ⁺); C ₂₆ H ₃₀ N ₃ SOBr requires 511 & 513. (DMSO) δ: 0.95 - 1.40 (7H,), 1.72 - 1.85 (4H, m), 2.42 (2H, m), 2.50 - 2.58 (4H, m, under DMSO), 2.87 - 2.95 (4H, m), 3.54 - 3.62 (1H, m), 6.48 (1H, d), 7.19 (1H, d, J = 5.4 Hz), 7.32 (1H, d), 7.48 (1H, d), 7.55 - 7.60 (2H, m), 7.72 (1H, d), 8.05 (1H, d).
66	7-CN	-C ₆ H ₄ (3-(2-pyridyl))	Found: 479 (MH ⁺); C ₃₁ H ₃₄ N ₄ O requires 478. (DMSO-d ₆) δ: 1.04 - 1.37 (2H, m), 1.28 - 1.47 (5H, m), 1.81 - 1.97 (4H, m), 2.49 (2H, t, J = 7 Hz), 2.56 - 2.61 (4H, m, obscured by DMSO), 2.92 - 3.00 (4H, m), 3.79 - 3.88 (1H, m), 7.38 (1H, d, J = 8 Hz), 7.44 - 7.47 (1H, m), 7.60 - 7.66 (3H, m), 7.90 - 8.00 (2H, m), 8.06 - 8.10 (1H, m), 8.25 - 8.27 (1H, m), 8.37 - 8.40 (1H, m), 8.55 (1H, s), 8.73 - 8.76 (1H, m).
67	7-CN	-C ₆ H ₄ (3-(5-pyrimidinyl))	Found: 480 (MH ⁺); C ₃₀ H ₃₃ N ₅ O requires 479. (DMSO-d ₆) & [HCl salt] δ: 1.05 - 1.12 (2H, m), 1.30 - 1.41 (3H, m), 1.65 - 1.70 (2H, m), 1.78 - 1.82 (2H, m), 1.88 - 1.92 (2H, m), 2.96 - 3.04 (2H, m), 3.08 - 3.20 (4H, m), 3.30 - 3.45 (2H, m), 3.65 - 3.71 (2H, m), 3.75 - 3.80 (1H, m), 7.45 (1H, d, J = 8 Hz), 7.60 - 7.73 (2H, m), 7.92 - 7.98 (3H, m), 8.23 (1H, s), 8.32 - 8.36 (1H, m), 9.21 - 9.23 (3H, m), 10.67 (1H, s).
68	7-CN	-C ₆ H ₄ (3-C ₆ H ₄ (4-CN))	Found: 503 (MH ⁺); C ₃₃ H ₃₄ N ₄ O requires 502. (DMSO-d ₆) δ: 1.00 - 1.11 (2H, m), 1.20 - 1.43 (5H, m), 1.79 - 1.83 (2H, m), 1.88 - 1.93 (2H, m), 2.48 (2H, t, J = 7 Hz), 2.52 - 2.58 (4H, m, obscured by DMSO), 2.91 -

			2.96 (4H, m), 3.75 - 8.83 (1H, m), 7.34 - (1H, d, J = 8 Hz), 7.58 - 7.61 (3H, m), 7.83 - 7.89 (3H, m), 7.95 - 8.05 (3H, m), 8.20 (1H, s), 8.33 - 8.35 (1H, m).
69	7-CN	-C ₆ H ₄ (3-(3-(5-ethyl)-1,2,4-oxadiazolyl))	Found: 498 (MH ⁺); C ₃₀ H ₃₅ N ₅ O ₂ requires 497. δ: 1.07 - 1.33 (5H, m), 1.41 - 1.50 (5H, m), 1.82 - 1.86 (2H, m), 2.09 - 2.13 (2H, m), 2.48 - 2.54 (2H, m), 2.61 - 2.64 (4H, m), 2.90 - 3.10 (6H, m), 3.89 - 4.04 (1H, m), 6.05 (1H, d, J = 8 Hz), 7.18 (1H, d, J = 8 Hz), 7.38 - 7.43 (2H, m), 7.57 (1H, t, J = 8 Hz), 7.99 (1H, dd, J = 8 Hz and 1 Hz), 8.20 (1H, dd, J = 8 Hz and 1 Hz), 8.33 (1H, d, J = 1 Hz).
70	7-CN	<i>trans</i> -CH=CH(2-thiophenyl)	Found: 434 (MH ⁺); C ₂₆ H ₃₁ N ₃ OS requires 433. (DMSOd ₆) δ: 0.85 - 1.30 (5H, m), 1.37 (2H, m), 1.90 (4H, m), 2.45 - 2.75 (6H, m), 3.00 (4H, m), 3.58 (1H, m), 6.35 (1H, d, J = 16 Hz), 7.10 (1H, m), 7.35 (2H, m), 7.45 - 7.65 (4H, m), 7.97 (1H, d, J = 16 Hz).
71	7-CN	<i>trans</i> -CH=CH(2-furanyl)	Found: 418 (MH ⁺); C ₂₆ H ₃₁ N ₃ O ₂ requires 417. (DMSOd ₆) δ: 0.80 - 1.30 (5H, m), 1.37 (2H, m), 1.78 (4H, m), 2.30 - 2.70 (6H, m), 2.93 (4H, m), 3.55 (1H, m), 6.38 (1H, d, J = 16 Hz), 6.55 (1H, dd, J = 3,2 Hz), 6.74 (1H, d, J = 3 Hz), 7.19 (1H, d, J = 16 Hz), 7.33 (1H, d, J = 8 Hz), 7.57 (2H, m), 7.75 (1H, s), 8.00 (1H, d, J = 8 Hz).
72	7-CN	<i>trans</i> -CH=CH(3-thiophenyl)	Found: 434 (MH ⁺); C ₂₆ H ₃₁ N ₃ OS requires 433. (DMSOd ₆) δ: 0.85 - 1.30 (5H, m), 1.40 (2H, m), 1.80 (4H, m), 2.35 - 2.70 (6H, m), 2.90 (4H, m), 3.60 (1H, m), 6.40 (1H, d, J = 16 Hz), 7.30 (2H, m), 7.38 (1H, d, J = 16

			Hz), 7.60 (3H, m), 7.75 (1H, m), 7.90 (1H, d, J = 8 Hz).
73	7-CN	<i>trans</i> -CH=CH(3-furanyl)	Found: 418 (MH ⁺); C ₂₆ H ₃₁ N ₃ O ₂ requires 417. (DMSOd ₆) δ: 0.85 - 1.30 (5H, m), 1.35 (2H, m), 1.80 (4H, m), 2.30 - 2.60 (6H, m), 2.85 (4H, m), 3.55 (1H, m), 6.28 (1H, d, J = 16 Hz), 6.66 (1H, s), 7.28 (1H, d, J = 16 Hz), 7.31 (1H, d, J = 8 Hz), 7.55 (2H, m), 7.71 (1H, s), 7.85 (1H, d, J = 8 Hz), 7.98 (1H, s).
74	7-CN	<i>trans</i> -CH=CH(4-quinoliny)	Found: 479 (MH ⁺); C ₃₁ H ₃₄ N ₄ O requires 478. δ (DMSO + TFA): 1.00 - 1.30 (5H, m), 1.60 - 1.70 (2H, m), 1.75 - 1.81 (2H, m), 1.90 - 1.95 (2H, m), 2.90 - 3.30 (8H, m), 3.60 - 3.80 (2H, m), 7.07 (1H, d, J = 16 Hz), 7.47 (1H, d, J = 8 Hz), 7.71 (1H, dd, J = 8 Hz), 7.75 (1H, s), 7.95 (1H, m), 8.05 - 8.30 (4H, m), 8.45 (1H, d, J = 8 Hz), 8.53 (1H, d, J = 8 Hz), 9.25 (1H, d, J = 5 Hz), 9.78 (1H, br s).
75		<i>trans</i> -CH=CH(5-pyrimidinyl)	Found: 430 (MH ⁺); C ₂₆ H ₃₁ N ₅ O requires 429. δ (DMSO + TFA): 1.00 - 1.30 (5H, m), 1.55 - 1.65 (2H, m), 1.75 - 1.80 (2H, m), 1.80 - 1.90 (2H, m), 3.00 - 3.25 (8H, m), 3.60 - 3.75 (3H, m), 6.80 (1H, d, J = 16 Hz), 7.42 (1H, d, J = 16 Hz), 7.46 (1H, d, J = 8 Hz), 7.71 (1H, dd, J = 8 Hz, 2 Hz), 7.74 (1H, s), 8.15 (1H, d, J = 8 Hz), 9.00 (2H, s), 9.15 (1H, s), 9.72 (1H, br s).
76	7-CN	-CH ₂ C ₆ H ₃ (2,4-diF)	Found: 452 (MH ⁺); C ₂₇ H ₃₁ F ₂ N ₃ O requires 451. δ (DMSO + TFA): 0.90 - 1.10 (2H, m), 1.20 - 1.30 (3H, m), 1.50 - 1.65 (2H, m), 1.70 - 1.85 (4H, m), 2.90 - 3.30 (8H, m),

			3.40 (2H, s), 3.50 (1H, m), 3.67 (2H, m), 7.00 (1H, m), 7.15 (1H, m), 7.40 (1H, m), 7.45 (1H, d, J = 8 Hz), 7.70 (1H, m), 7.73 (1H, s), 7.96 (1H, d, J = 8 Hz), 9.70 (1H, br s).
77	7-CN	-CH ₂ (1-naphthyl)	Found: 466 (MH ⁺); C ₃₁ H ₃₅ N ₃ O requires 465. δ: 0.70 - 0.80 (2H, m), 0.90 - 1.10 (3H, m), 1.30 - 1.40 (2H, m), 1.60 - 1.70 (2H, m), 1.70 - 1.80 (2H, m), 2.40 (2H, m), 2.55 (4H, m), 2.80 - 3.00 (4H, m), 3.66 (1H, m), 4.00 (2H, s), 5.05 (1H, d, J = 8 Hz), 7.15 (1H, d, J = 8 Hz), 7.34 (1H, s), 7.35 - 7.40 (2H, m), 7.45 (1H, m), 7.50 (2H, m), 7.83 (1H, d, J = 8 Hz), 7.86 (1H, m), 7.93 (1H, m).
78	7-COMe	3-pyrrolo[2,3-b]pyridyl	Found: 459 (MH ⁺); C ₂₈ H ₃₄ N ₄ O ₂ requires 458. (DMSOd ₆) δ: 1.0 - 1.20 (2H, m), 1.25 - 1.55 (5H, m), 1.75 - 2.00 (4H, m), 2.40 - 2.65 (6H, m), 2.56 (3H, s), 2.95 (4H, m), 3.75 (1H, m), 7.15 (1H, m), 7.25 (1H, m), 7.74 (3H, m), 8.15 (1H, s), 8.25 (1H, m), 8.45 (1H, m), 12.05 (1H, br s).
79	7-COMe	-CH ₂ C ₆ H ₄ (4-F)	Found: 451 (MH ⁺); C ₂₈ H ₃₅ FN ₂ O ₂ requires 450. (DMSOd ₆) δ: 0.85 - 1.20 (5H, m), 1.35 (2H, m), 1.85 (4H, m), 2.40 - 2.65 (6H, m), 2.54 (3H, s), 2.76 (4H, m), 3.35 (2H, s), 3.45 (1H, m), 7.10 (2H, m), 7.25 (3H, m), 7.65 (2H, m), 7.90 (1H, d, J = 8 Hz).
80	7-COMe	-C ₆ H ₄ (3-(3-(5-methyl)-1,2,4-oxadiazolyl))	Found: 501 (MH ⁺); C ₃₀ H ₃₆ N ₄ O ₃ requires 500. (DMSOd ₆) δ: 0.90 - 1.45 (7H, m), 1.84 (4H, m), 2.40 - 2.60 (6H, m), 2.55 (3H, s), 2.70 (3H, s), 2.94 (4H, m), 3.91 (1H, m), 7.26 (1H, d, J = 8 Hz), 7.60 - 7.75 (3H, m),

			8.05 (1H, m), 8.10 (1H, m), 8.45 (2H, m).
81	7-COMe	<i>trans</i> -CH=CHC ₆ H ₄ (4-F)	Found: 463 (MH ⁺); C ₂₉ H ₃₅ FN ₂ O ₂ requires 462. δ: (DMSOd ₆) δ: 0.95 - 1.30 (5H, m), 1.40 (2H, m), 1.82 (4H, m), 2.40 - 2.65 (6H, m), 2.56 (3H, s), 2.95 (4H, m), 3.62 (1H, m), 6.56 (1H, d, J = 16 Hz), 7.25 (3H, m), 7.40 (1H, d, J = 16 Hz), 7.65 (2H, m), 7.70 (2H, m), 7.95 (1H, d, J = 8 Hz).
82	7-CN	-CH ₂ (6-(2-amino)benzothiazolyl)	Found: 488 (MH ⁺); C ₂₈ H ₃₃ N ₅ OS requires 487. δ: 0.85 - 1.11 (5H, m), 1.37 (2H, m), 1.68 (2H, m), 1.89 (2H, m), 2.45 (2H, m), 2.59 (4H, m), 2.92 (4H, m), 3.57 (2H, s), 3.68 (1H, m), 5.16 (3H, m), 7.17 (2H, m), 7.38 (2H, m), 7.52 (2H, m).
83	7-CN	-CH ₂ (6-(2-methyl)benzothiazolyl)	Found: 487 (MH ⁺); C ₂₉ H ₃₄ N ₄ OS requires 486. δ (DMSO-d ₆): 0.93 (2H, m), 1.15 (3H, m), 1.33 (2H, m), 1.75 (4H, m), 2.42 (2H, m), 2.52 (4H, m), 2.77 (3H, s), 2.90 (4H, m), 3.43 (3H, m), 7.33 (2H, m), 7.56 (2H, m), 7.82 (2H, m), 7.95 (1H, d, J = 8 Hz).
84	7-CN	-CH ₂ (6-(2,3-dihydro-2-oxo)indolinylyl)	Found: 471 (MH ⁺); C ₂₉ H ₃₄ N ₄ O ₂ requires 470. δ (DMSO-d ₆ + TFA): 0.88 - 1.32 (5H, m), 1.59 (2H, m), 1.74 (4H, m), 2.90 - 3.27 (8H, m), 3.29 (2H, s), 3.40 (2H, s), 3.45 (1H, m), 3.65 (2H, m), 6.73 (1H, s), 6.79 (1H, d, J = 9 Hz), 7.08 (1H, d, J = 9 Hz), 7.45 (1H, d, J = 9 Hz), 7.69 (1H, d, J = 9 Hz), 7.72 (1H, s), 7.90 (1H, d, J = 9 Hz), 9.86 (1H, br s), 10.33 (1H, br s).
85	7-CN	-CH ₂ (5-(2,3-dihydro-2-oxo)indolinylyl)	Found: 471 (MH ⁺); C ₂₉ H ₃₄ N ₄ O ₂ requires 470. δ (DMSO-d ₆ + TFA): 0.90 - 1.35 (5H, m),

			1.59 (2H, m), 1.75 (4H, m), 2.91 - 3.29 (8H, m), 3.27 (2H, s), 3.43 (2H, s), 3.47 (1H, m), 3.66 (2H, m), 6.72 (1H, d, J = 9 Hz), 7.01 (1H, d, J = 9 Hz), 7.06 (1H, s), 7.45 (1H, d, J = 9 Hz), 7.69 (1H, d, J = 9 Hz), 7.72 (1H, s), 7.88 (1H, d, J = 9 Hz), 9.90 (1H, br s), 10.29 (1H, br s).
86	7-CN	CH=CHC ₆ H ₄ (4-CONHMe)	Found: 485 (MH ⁺); C ₃₀ H ₃₆ N ₄ O ₂ requires 484. (DMSO + TFA) δ: 0.97 - 1.28 (5H, m), 1.58 - 1.64 (2H, m), 1.76 - 1.90 (4H, m), 2.79 (3H, m), 2.99 - 3.33 (8H, m), 3.66 - 3.77 (3H, m), 6.69 (1H, d, J = 16 Hz), 7.41 - 7.45 (2H, m), 7.60 - 7.73 (3H, m), 7.86 (2H, d, J = 8 Hz), 8.03 (1H, d), 8.45 (1H, m), 9.75 (1H, b s).
87	7-CN	CH ₂ (5-(2-amino)benzoxazolyl)	Found: 472 (MH ⁺); C ₂₈ H ₃₃ N ₅ O ₂ requires 471. (DMSO + TFA) δ: 0.77 - 1.23 (5H, m), 1.54 - 1.62 (2H, m), 1.70 - 1.79 (4H, m), 2.99 - 3.24 (8H, m), 3.38 - 3.45 (3H, m), 3.60 - 3.69 (2H, m), 6.49 (1H, d, J = 10 Hz), 7.15 (1H, s), 7.32 (1H, d, J = 8 Hz), 7.45 (1H, d, J = 8 Hz), 7.67 - 7.72 (2H, m), 7.93 (1H, d), 8.21 (2H, b s), 9.92 (1H, bs).
88	7-CN	-CH ₂ (6-(1,2-dihydro-2-oxo)quinoliny)	Found: 483 (MH ⁺); C ₃₀ H ₃₄ N ₄ O ₂ requires 482. δ (DMSO + TFA): 0.90 - 1.20 (5H, m), 1.50 - 1.65 (2H, m), 1.70 - 1.80 (4H, m), 2.90 - 3.30 (8H, m), 3.40 (2H, s), 3.45 (1H, m), 3.60 - 3.70 (2H, m), 6.48 (1H, d, J = 10 Hz), 7.22 (1H, d, J = 8 Hz), 7.50 (1H, dd, J = 10 Hz, 2 Hz), 7.60 - 7.50 (2H, m), 7.80 (1H, dd, J = 10 Hz, 2 Hz), 7.70 (1H, s), 7.86 (1H, d, J = 9 Hz), 7.95 (1H, d, J = 8 Hz), 9.80 (1H, br s), 11.70 (1H, br s).
89	7-CN	-CH ₂ (7-(1,2-	Found: 483 (MH ⁺); C ₃₀ H ₃₄ N ₄ O ₂

		dihydro-2-oxoquinolinyI)	requires 482. δ (DMSO): 0.90 - 1.00 (2H, m), 1.10 - 1.20 (5H, m), 1.25 - 1.40 (2H, m), 1.70 - 1.80 (4H, m), 2.40 - 2.50 (4H, m), 2.80 - 2.90 (4H, m), 3.30 - 3.45 (3H, m), 6.43 (1H, d, J = 8 Hz), 7.05 (1H, d, J = 8 Hz), 7.17 (1H, s), 7.32 (1H, d, J = 8 Hz), 7.50 - 7.60 (3H, m), 7.84 (1H, d, J = 9 Hz), 7.97 (1H, d, J = 8 Hz), 11.70 (1H, s).
90	7-COMe	<i>trans</i> -CH=CHC ₆ H ₄ (3-OMe)	Found: 475 (MH ⁺); C ₃₀ H ₃₈ N ₂ O ₃ requires 474. δ : 1.05 - 1.40 (5H, m), 1.44 (2H, m), 1.81 (2H, m), 2.05 (2H, m), 2.48 (2H, m), 2.58 (3H, s), 2.62 (4H, m), 2.98 (4H, m), 3.82 (3H, s), 3.85 (1H, m), 5.43 (1H, d, J = 8 Hz), 6.33 (1H, d, J = 16 Hz), 6.89 (1H, dd, J = 2, 8 Hz), 7.00 (1H, m), 7.08 (1H, d, J = 8 Hz), 7.17 (1H, d, J = 8 Hz), 7.29 (1H, m), 7.56 (1H, d, J = 16 Hz), 7.70 (1H, s), 7.73 (1H, m).
91	7-COMe	<i>trans</i> -CH=CHC ₆ H ₄ (2-CN)	Found: 470 (MH ⁺); C ₃₀ H ₃₅ N ₃ O ₂ requires 469. δ : 1.05 - 1.35 (5H,), 1.43 (2H, m), 1.81 (2H, m), 2.06 (2H, m), 2.49 (2H, m), 2.58 (3H, s), 2.63 (4H, m), 2.98 (4H, m), 3.86 (1H, m), 5.62 (1H, d, J = 8 Hz), 6.66 (1H, d, J = 16 Hz), 7.17 (1H, d, J = 8 Hz), 7.43 (1H, m), 7.60 (2H, m), 7.70 (3H, m), 7.77 (1H, d, J = 16 Hz).
92	7-COMe	<i>trans</i> -CH=CH(3-thiophenyl)	Found: 451 (MH ⁺); C ₂₇ H ₃₄ N ₂ O ₂ S requires 450. δ : 1.05 - 1.35 (5H, m), 1.42 (2H, m), 1.80 (2H, m), 2.05 (2H, m), 2.49 (2H, m), 2.58 (3H, s), 2.62 (4H, m), 2.98 (4H, m), 3.85 (1H, m), 5.41 (1H, d, J = 8 Hz), 6.18 (1H, d, J = 16 Hz), 7.15 (1H, d, J = 8 Hz), 7.25 (2H, m), 7.42 (1H, m), 7.59 (1H, d, J = 16

			Hz), 7.69 (1H, s), 7.72 (1H, m).
93	7-COMe	<i>trans</i> -CH=CH(8-(1,2-dihydro-2-oxo)-quinolinyl)	Found: 512 (MH ⁺); C ₃₂ H ₃₇ N ₃ O ₃ requires 511. DMSOd ₆ , HCl salt) δ : 1.00 - 1.40 (5H, m), 1.60 - 1.95 (6H, m), 2.56 (3H, s), 2.90 - 3.20 (6H, m), 3.40 (2H, m), 3.65 (3H, m), 6.55 (2H, m), 7.22 (1H, t, J = 8 Hz), 7.37 (1H, d, J = 8 Hz), 7.70 (2H, m), 7.80 (2H, m), 7.90 - 8.10 (3H, m), 10.59 (1H, br s), 11.35 (1H, br s).
94	7-CN	-C ₆ H ₄ (3-(1-pyrazolyl))	Found: 468 (MH ⁺); C ₂₉ H ₃₃ N ₅ O requires 467. δ : 1.06 - 1.40 (5H, m), 1.40 - 1.50 (2H, m), 1.81 - 1.86 (2H, m), 2.08 - 2.13 (2H, m), 2.38 - 2.54 (2H, m), 2.61 - 2.65 (4H, m), 2.90 - 3.00 (4H, m), 3.89 - 3.96 (1H, m), 6.06 (1H, d, J = 8 Hz), 6.50 (1H, t, J = 2 Hz), 7.18 (1H, d, J = 8 Hz), 7.29 - 7.43 (2H, m), 7.51 (1H, t, J = 8 Hz), 7.68 (1H, d, J = 8 Hz), 7.75 (1H, d, J = 1.5 Hz), 7.79 - 7.83 (1H, m), 8.00 (1H, m), 8.08 (1H, m).
95	7-CN	-CH ₂ (2-thiophenyl)	Found: 422 (MH ⁺); C ₂₅ H ₃₁ N ₃ OS requires 421. δ : 0.90 - 1.20 (5H, m), 1.35 - 1.46 (2H, m), 1.69 - 1.72 (2H, m), 1.80 - 2.00 (2H, m), 2.42 - 2.48 (2H, m), 2.57 - 2.65 (4H, m), 2.90 - 2.96 (4H, m), 3.65 - 3.75 (1H, m), 3.74 (2H, s), 5.38 (1H, d, J = 8 Hz), 8.92 (1H, m), 6.98 (1H, m), 7.16 (1H, d, J = 8 Hz), 7.23 - 7.26 (1H, m), 7.36 - 7.42 (2H, m).
96	7-CN	-CH ₂ (3-benzothiophenyl)	Found: 472 (MH ⁺); C ₂₉ H ₃₃ N ₃ OS requires 471. δ : 0.80 - 1.20 (5H, m), 1.30 - 1.40 (2H, m), 1.60 - 1.75 (2H, m), 1.80 - 1.90 (2H, m), 2.40 - 2.50 (2H, m), 2.50 - 2.70 (4H, m), 2.85 - 2.95 (4H, m), 3.65 - 3.75 (1H, m),


			3.80 (2H, s), 5.23 (1H, d, J = 8 Hz), 7.16 (1H, d, J = 8 Hz), 7.30 - 7.45 (5H, m), 7.60 - 7.70 (1H, m), 7.85 - 7.92 (1H, m).
97	7-CN	-C ₆ H ₄ (3-(2-(5-methyl)-1,3,4-oxadiazolyl))	Found: 484 (MH ⁺); C ₂₉ H ₃₃ N ₅ O ₂ requires 483. δ: 1.10 - 1.40 (5H, m), 1.41 - 1.50 (2H, m), 1.82 - 1.87 (2H, m), 2.10 - 2.14 (2H, m), 2.48 - 2.54 (2H, m), 2.62 - 2.65 (7H, m), 2.93 - 3.00 (4H, m), 3.93 - 3.97 (1H, m), 6.04 (1H, d, J = 8 Hz), 7.18 (1H, d, J = 8 Hz), 7.35 - 7.43 (2H, m), 7.59 (1H, t, J = 8 Hz), 7.97 (1H, dd, J = 6 and 1 Hz), 8.11 (1H, dd, J = 8 and 1 Hz), 8.36 (1H, d, J = 1 Hz).
98	7-CN	<i>trans</i> -CH=CH(2-naphthyl)	Found: 478 (MH ⁺); C ₃₂ H ₃₅ N ₃ O requires 477. δ (DMSO + TFA): 1.00 - 1.65 (5H, m), 1.63 (2H, m), 1.76 - 1.91 (4H, m), 2.97 - 3.32 (8H, m), 3.66 - 3.93 (3H, m), 6.75 (1H, d, J = 16 Hz), 7.47 (1H, d), 7.54 - 7.62 (3H, m), 7.65 - 7.80 (3H, m), 7.85 - 7.95 (3H, m), 8.00 - 8.10 (2H, m), 9.77 (1H, b s).
99	7-COMe	CH ₂ (3-benzothiophenyl)	Found: 489 (MH ⁺); C ₃₀ H ₃₆ N ₂ O ₂ S requires 488. δ: 0.80 - 1.25 (5H, m), 1.37 (2H, m), 1.70 (2H, m), 1.85 (2H, m), 2.41 (2H, m), 2.57 (3H, s), 2.59 (4H, m), 2.94 (4H, m), 3.67 (1H, m), 3.80 (2H, s), 5.23 (1H, d, J = 8 Hz), 7.16 (1H, d, J = 8 Hz), 7.32 (1H, s), 7.40 (2H, m), 7.67 (1H, s), 7.70 (2H, m), 7.88 (1H, m).
100	7-COMe	<i>trans</i> -CH=CC ₆ H ₄ (4-NHCOMe)	Found: 502 (MH ⁺); C ₃₁ H ₃₉ N ₃ O ₃ requires 501. δ: 0.90 - 1.35 (5H, m), 1.39 (2H, m), 1.80 (4H, m), 2.05 (3H, s), 2.50 - 2.80 (6H, m), 2.54 (3H, s), 2.94 (4H, m), 3.60 (1H, m),

			6.47 (1H, d, J = 16 Hz), 7.20 - 7.40 (2H, m), 7.46 (2H, d, J = 9 Hz), 7.61 (2H, d, J = 9 Hz), 7.71 (2H, m), 7.91 (1H, d, J = 8 Hz), 10.09 (1H, s).
101	7-COMe	-CH ₂ (6-(2-amino)-benzothiazolyl)	Found: 505 (MH ⁺); C ₂₉ H ₃₆ N ₄ O ₂ S requires 504. δ (DMSOd ₆): 0.85 - 1.30 (5H, m), 1.48 (2H, m), 1.75 (4H, m), 2.55 (3H, s), 2.70 - 3.25 (10H, m), 3.34 (2H, s, obscured by H ₂ O), 3.55 (1H, m), 7.05 (1H, m), 7.15 - 7.35 (2H, m), 7.39 (2H, br s), 7.47 (1H, m), 7.75 (2H, m), 7.90 (1H, d, J = 8 Hz).
102	7-COMe	8-(1,4-dihydro-4-oxo)-quinolinyl	Found: 486 (MH ⁺); C ₃₀ H ₃₅ N ₃ O ₃ required 485. δ (DMSOd ₆): 0.95 - 1.20 (2H, m), 1.20 - 1.70 (5H, m), 1.70 - 2.00 (4H, m), 2.56 (3H, s), 2.65 - 3.20 (10H, m), 3.82 (1H, m), 6.08 (1H, d, J = 7 Hz), 7.25 - 7.45 (2H, m), 7.77 (2H, m), 7.93 (1H, m), 8.10 (1H, m), 8.26 (1H, d, J = 8 Hz), 8.71 (1H, d, J = 8 Hz), 12.05 (1H, m).
103	7-COMe	<i>trans</i> -CH=CHC ₆ H ₄ (2-COMe)	Found: 487 (MH ⁺); C ₃₁ H ₃₈ N ₂ O ₃ requires 486. δ: 1.05 - 1.40 (5H, m), 1.44 (2H, m), 1.80 (2H, m), 2.05 (2H, m), 2.50 (2H, m), 2.58 (3H, s), 2.60 (3H, s), 2.64 (4H, m), 2.98 (4H, m), 3.85 (1H, m), 5.50 (1H, d, J = 8 Hz), 6.20 (1H, d, J = 16 Hz), 7.18 (1H, d, J = 8 Hz), 7.35 - 7.65 (3H, m), 7.69 (3H, m), 7.91 (1H, d, J = 16 Hz).
104	7-COMe	-CH ₂ (2-benzothiophenyl)	Found: 489 (MH ⁺); C ₃₀ H ₃₆ N ₂ O ₂ S requires 488. δ: 1.00 - 1.30 (5H, m), 1.40 (2H, m), 1.72 (2H, m), 1.94 (2H, m), 2.44 (2H, m), 2.57 (3H, s), 2.61 (4H, m), 2.95 (4H, m), 3.73 (1H, m), 3.82 (2H, s), 5.47 (1H, d, J = 8 Hz), 7.16 (2H, m), 7.30 - 7.39 (2H, m),

			7.65 - 7.81 (4H, m).
105	7-CN	<i>trans</i> -CH=CH(5-(3-acetyl)indolyl)	<p>Found: 509 (MH⁺); C₃₂H₃₆N₄O₂ requires 508.</p> <p>δ (DMSOd₆): 0.85 - 1.28 (5H, m), 1.35 (2H, m), 1.65 - 1.95 (4H, m), 2.35 - 2.65 (6H, m), 2.46 (3H, s), 2.91 (4H, m), 3.59 (1H, m), 6.60 (1H, d, J = 16 Hz), 7.30 - 7.65 (6H, m), 7.96 (1H, d, J = 8 Hz), 8.37 (2H, m), 12.05 (1H, br s).</p>
106	7-CN	-C ₆ H ₄ (5-(3-methyl)-1,2,4-oxadiazolyl)	<p>Found: 484 (MH⁺); C₂₉H₃₃N₅O₂ requires 483.</p> <p>δ (CDCl₃): 1.13 - 1.28 (5H, m), 1.43 - 1.48 (2H, m), 1.83 - 1.86 (2H, m), 2.10 - 2.13 (2H, m), 2.49 (3H, s), 2.51 (2H, m), 2.62 - 2.64 (4H, m), 2.88 - 2.98 (4H, m), 3.94 - 3.98 (1H, m), 6.02 (1H, d, J = 8 Hz), 7.18 (1H, d, J = 7.7 Hz), 7.38 (1H, s), 7.39 (1H, d, J = 7.7 Hz), 7.64 (1H, t, J = 7.8 Hz), 8.05 (1H, d), 8.21 (1H, d), 8.39 (1H, br s).</p>
107	7-CN	-CH ₂ (5-(2-methyl)-benzimidazolyl)	<p>Found: 470 (MH⁺); C₂₉H₃₅N₅O requires 469.</p> <p>δ: 0.87 - 1.09 (5H, m), 1.14 (1H, br s), 1.37 (2H, m), 1.70 (2H, m), 1.88 (2H, m), 2.45 (2H, m), 2.56 (3H, s), 2.60 (4H, m), 2.93 (4H, m), 3.61 (2H, s), 3.69 (1H, m), 3.61 (2H, s), 3.69 (1H, m), 5.44 (1H, d, J = 7 Hz), 7.04 (1H, dd, J = 8, 2 Hz), 7.15 (1H, d, J = 8 Hz), 7.30 - 7.47 (4H, m).</p>
108	7-CN	-CH ₂ (6-quinoxalinylyl)	<p>Found: 468 (MH⁺); C₂₉H₃₃N₅O requires 467.</p> <p>δ: 0.90 - 1.15 (5H, m), 1.40 (2H, m), 1.73 (2H, m), 1.93 (2H, m), 2.45 (2H, m), 2.59 (4H, m), 2.94 (4H, m), 3.67 (1H, m), 3.76 (2H, s), 5.33 (1H, d, J = 7 Hz), 7.16 (1H, d, J = 8 Hz), 7.31 - 7.44 (2H, m), 7.72 (1H, dd, J = 9, 2 Hz), 7.96 (1H, d, J = 2 Hz),</p>

			8.08 (1H, d, J = 9 Hz), 8.85 (2H, s).
109	7-CN	<i>trans</i> -CH=CH(3-(2-acetyl)furanyl)	<p>Found: 460 (MH⁺); C₂₈H₃₃N₃O₃ requires 459.</p> <p>δ: 1.05 - 1.35 (5H, m), 1.45 (2H, m), 1.80 (2H, m), 2.04 (2H, m), 2.48 (2H, m), 2.53 (3H, s), 2.61 (4H, m), 2.95 (4H, m), 3.84 (1H, m), 5.56 (1H, d, J = 8 Hz), 6.43 (1H, d, J = 16 Hz), 6.70 (1H, d, J = 2 Hz), 7.17 (1H, d, J = 8 Hz), 7.38 (1H, s), 7.41 (1H, d, J = 8 Hz), 7.45 (1H, d, J = 2 Hz), 7.95 (1H, d, J = 16 Hz).</p>
110	7-CN	-CH ₂ (6-(2-amino)benzoxazolyl)	<p>Found: 472 (MH⁺); C₂₈H₃₃N₅O₂ requires 471.</p> <p>δ: 0.81 - 1.12 (5H, m), 1.40 (2H, m), 1.72 (2H, m), 1.89 (2H, m), 2.45 (2H, m), 2.93 (4H, m), 3.58 (2H, s), 3.69 (1H, m), 4.92 (2H, br s), 5.13 (1H, m), 7.05 (1H, m), 7.17 (2H, m), 7.36 (3H, m).</p>
111	7-CN	-CH ₂ (6-(3,4-dihydro-2-oxo)-2H-benzoxazinyl)	<p>Found: 487 (MH⁺); C₂₉H₃₄N₄O₃ requires 486.</p> <p>δ (DMSO-d₆): 0.83 - 1.24 (5H, m), 1.32 (2H, m), 1.73 (4H, m), 2.47 (6H, m), 2.90 (4H, m), 3.25 (2H, s), 3.40 (1H, m), 4.53 (2H, s), 6.80 (3H, m), 7.31 (1H, d, J = 8 Hz), 7.56 (2H, m), 7.84 (1H, d, J = 8 Hz), 10.62 (1H, br s).</p>
112	7-CN	<i>trans</i> -CH=CHC ₆ H ₃ (2-F, 5-NHCOMe)	<p>Found: 503 (MH⁺); C₃₀H₃₅FN₄O₂ requires 502.</p> <p>δ (DMSO-d₆ + TFA): 0.94 - 1.34 (5H, m), 1.61 (2H, m), 1.76 (2H, m), 1.87 (2H, m), 2.05 (3H, s), 2.93 - 3.33 (8H, m), 3.54 - 3.77 (3H, m), 6.65 (1H, d, J = 15 Hz), 7.20 (1H, t, J = 9 Hz), 7.43 (3H, m), 7.69 (1H, d, J = 9 Hz), 7.73 (1H, s), 8.03 (1H, m), 8.19 (1H, d, J = 9 Hz), 9.88 (1H, br s), 10.09 (1H, br s).</p>

Table 2.



Example No.	R	Mass spectrum, ¹ H NMR
113	-CH ₂ -(2-benzothiophenyl)	<p>Mass spectrum (API⁺): Found 529 (MH⁺). C₃₁H₃₆N₄O₂S requires 528.</p> <p>NMR (CDCl₃) δ: 1.00 - 1.10 (4H, m), 1.19 (1H, m), 1.35 - 1.45 (2H, m), 1.75 (2H, m), 1.95 (2H, m), 2.40 - 2.50 (5H, m), 2.61 (4H, m), 2.97 (4H, m), 3.73 (1H, m), 3.82 (2H, s), 5.46 (1H, d, J = 8 Hz), 7.16 (1H, s), 7.22 (1H, d, J = 8 Hz), 7.25 - 7.42 (2H, m), 7.73 (1H, d, J = 8 Hz), 7.79 (1H, d, J = 8 Hz), 7.81 - 7.88 (2H, m).</p>
114	(E)-CH=CH-(3-thienyl)	<p>Mass spectrum (API⁺): Found 491 (MH⁺). C₂₈H₃₄N₄O₂S requires 490.</p> <p>NMR (CDCl₃) δ: 1.04 - 1.15 (4H, m), 1.25 (1H, m), 1.44 (2H, m), 1.76 (2H, m), 2.05 (2H, m), 2.46 (3H, s), 2.50 (2H, m), 2.64 (4H, m), 3.00 (4H, m), 3.85 (1H, m), 5.36 (1H, d, J = 8 Hz), 6.18 (1H, d, J = 16 Hz), 7.22 - 7.20 (2H, m), 7.30 (1H, m), 7.43 (1H, m), 7.59 (1H, d, J = 16 Hz), 7.80 - 7.90 (2H, m).</p>
115	5-quinolyl	<p>Mass spectrum (API⁺): Found 510 (MH⁺). C₃₁H₃₅N₅O₂ requires 509.</p> <p>NMR (CDCl₃) δ: 1.15 - 1.27 (5H, m), 1.45 (2H, m), 1.85 (2H, m), 2.20 (2H, m), 2.46 (3H, s), 2.55 (2H, m), 2.70 (4H, m), 3.00 (4H, m), 4.00 (1H, m), 5.85 (1H, d, J = 8 Hz), 7.25 (1H, d, J = 8 Hz), 7.46 (1H, dd, J = 4, 8 Hz), 7.60 - 7.72 (2H, m), 7.84 - 7.87 (2H, m), 8.16 (1H, d, J = 8 Hz), 8.72 (1H, d, J = 8 Hz), 8.92 (1H, m).</p>
116	3- pyrrolo[2,3-b]pyridyl	<p>Mass spectrum (API⁺): Found 499 (MH⁺). C₂₉H₂₄N₆O requires 498.</p>


		NMR (DMSO-d ₆) δ : 0.90 - 1.10 (2H, m), 1.10 - 1.40 (5H, m), 1.70 - 1.90 (4H, m), 2.40 - 2.70 (6H, m), 2.96 (3H, s), 3.31 (4H, m), 3.89 (1H, m), 7.15 (1H, m), 7.36 (1H, d, J = 8 Hz), 7.71 (1H, d, J = 8 Hz), 7.75 - 7.85 (2H, m), 8.12 (1H, s), 8.20 (1H, s), 8.35 (1H, d, J = 8 Hz), 12.02 (1H, br s).
117	3-(3-(5-methyl)-1,2,4-oxadiazolyl)phenyl	Mass spectrum (API ⁺): Found 541 (MH ⁺). C ₃₁ H ₃₆ N ₆ O ₃ requires 540. NMR (CDCl ₃) δ 1.10 - 1.22 (4H, m), 1.27 (1H, m), 1.55 (2H, m), 1.90 (2H, m), 2.10 (2H, m), 2.47 (3H, s), 3.65 (2H, m), 2.68 (3H, s), 2.76 (4H, m), 3.06 (4H, m), 3.95 (1H, m), 6.00 (1H, d, J = 8 Hz), 7.25 (1H, d, J = 8 Hz), 7.57 (1H, t, J = 8 Hz), 7.80 - 7.90 (2H, m), 8.02 (1H, d, J = 8 Hz), 8.15 (1H, d, J = 8 Hz), 8.32 (1H, s).
118	8-(1,4-dihydro-4-oxo)quinolyl	Mass spectrum (API ⁺): Found 526 (MH ⁺). C ₃₁ H ₃₅ N ₅ O ₃ requires 525. NMR (DMSO-d ₆) δ : 0.90 - 1.10 (2H, m), 1.20 - 1.40 (5H, m), 1.80 - 2.00 (4H, m), 2.30 - 2.75 (9H, m), 2.96 (4H, m), 3.80 (1H, m), 6.09 (1H, d, J = 8 Hz), 7.30 - 7.40 (2H, m), 7.75 - 7.88 (2H, m), 7.92 (1H, m), 8.05 (1H, d, J = 8 Hz), 8.22 (1H, d, J = 8 Hz), 8.65 (1H, d, J = 8 Hz), 12.04 (1H, br s).
119	(E)-CH=CH-(4-fluoro)phenyl	Mass spectrum (API ⁺): Found 503 (MH ⁺). C ₃₀ H ₃₅ FN ₄ O ₂ requires 502. NMR (CDCl ₃) δ : 1.10 - 1.30 (5H, m), 1.40 - 1.47 (2H, m), 1.78 - 1.82 (2H, m), 2.00 - 2.10 (2H, m), 2.46 (3H, s), 2.47 - 2.52 (2H, m), 2.60 - 2.70 (4H, m), 2.95 - 3.05 (4H, m), 3.86 (1H, m), 5.38 (1H, d, J = 8 Hz), 6.26 (1H, d, J = 16 Hz), 7.05 (2H, t, J = 8 Hz), 7.24 (1H, d, J = 8 Hz), 7.47 (2H, dd, J = 5, 8 Hz), 7.57 (1H, d, J = 16 Hz), 7.80 - 7.90 (2H, m).
120	(E)-CH=CH-(3-fluoro)phenyl	Mass spectrum (API ⁺): Found 503 (MH ⁺). C ₃₀ H ₃₅ FN ₄ O ₂ requires 502.

		NMR (CDCl ₃) δ : 1.10 - 1.30 (5H, m), 1.42 (2H, m), 1.81 (2H, m), 2.06 (2H, m), 2.46 (3H, s), 2.51 (2H, m), 2.65 (4H, m), 3.00 (4H, m), 3.87 (1H, m), 5.41 (1H, d, J = 8 Hz), 6.33 (1H, d, J = 16 Hz), 7.02 (1H, m), 7.15 (1H, m), 7.25 (2H, m), 7.31 (1H, m), 7.57 (1H, d, J = 16 Hz), 7.80 - 7.90 (2H, m).
121	(<i>E</i>)-CH=CH-(3-acetamido-2-fluoro)phenyl	Mass spectrum (API ⁺): Found 560 (MH ⁺). C ₃₂ H ₃₈ FN ₅ O ₃ requires 559. NMR (CDCl ₃) δ : 1.10 - 1.20 (4H, m), 1.20 - 1.30 (1H, m), 1.40 - 1.50 (2H, m), 1.77 - 1.83 (2H, m), 2.05 - 2.12 (2H, m), 2.24 (3H, s), 2.46 (3H, s), 2.55 (2H, m), 2.65 (4H, m), 3.00 (4H, m), 3.85 (1H, m), 5.42 (1H, d, J = 8 Hz), 6.42 (1H, d, J = 16 Hz), 7.12 (1H, t, J = 8 Hz), 7.18 - 7.30 (2H, m), 7.38 (1H, s), 7.71 (1H, d, J = 16 Hz), 7.80 - 7.90 (2H, m), 8.30 (1H, m).
122	(<i>E</i>)-CH=CH-(3-acetyl)phenyl	Mass spectrum (API ⁺): Found 527 (MH ⁺). C ₃₂ H ₃₈ N ₄ O ₃ requires 526. NMR (CDCl ₃) δ : 1.10 - 1.20 (4H, m), 1.20 - 1.30 (1H, m), 1.50 (2H, m), 1.80 (2H, m), 2.05 (2H, m), 2.46 (3H, s), 2.56 (2H, m), 2.60 (3H, s), 2.65 (4H, m), 3.00 (4H, m), 3.85 (1H, m), 5.48 (1H, d, J = 8 Hz), 6.20 (1H, d, J = 16 Hz), 7.24 (1H, d, J = 8 Hz), 7.40 (1H, m), 7.45 - 7.55 (2H, m), 7.70 (1H, d, J = 8 Hz), 7.85 - 7.88 (2H, m), 7.91 (1H, d, J = 16 Hz).
123	-CH ₂ -(3-fluoro)phenyl	Mass spectrum (API ⁺): Found 491 (MH ⁺). C ₂₉ H ₃₅ FN ₄ O ₂ requires 490. NMR (CDCl ₃) δ : 1.00 - 1.12 (4H, m), 1.19 (1H, m), 1.40 (2H, m), 1.75 (2H, m), 1.93 (2H, m), 2.40 - 2.50 (5H, m), 2.62 (4H, m), 2.95 (4H, m), 3.52 (2H, s), 3.70 (1H, m), 5.14 (1H, d, J = 8 Hz), 6.90 - 7.05 (3H, m), 7.22 (1H, d, J = 8 Hz), 7.30 (1H, m), 7.80 - 7.90 (2H, m).
124	-CH ₂ -(2.4-	Mass spectrum (API ⁺): Found 509 (MH ⁺).

	difluoro)phenyl	<p>$C_{29}H_{34}F_2N_4O_2$ requires 508.</p> <p>NMR ($CDCl_3$) δ: 1.00 - 1.10 (4H, m), 1.15 - 1.25 (1H, m), 1.35 - 1.45 (2H, m), 1.70 - 1.80 (2H, m), 1.90 - 2.00 (2H, m), 2.46 (3H, s), 2.48 (2H, m), 2.63 (4H, m), 2.97 (4H, m), 3.48 (2H, s), 3.70 (1H, m), 5.24 (1H, d, $J = 8$ Hz), 6.85 (2H, m), 7.23 (1H, d, $J = 8$ Hz), 7.24 - 7.35 (1H, m), 7.80 - 7.90 (2H, m).</p>
125	2-naphthyl	<p>Mass spectrum (API⁺): Found 509 (MH⁺). $C_{32}H_{36}N_4O_2$ requires 508.</p> <p>NMR ($CDCl_3$) δ: 1.10 - 1.35 (5H, m), 1.40 - 1.50 (2H, m), 1.80 - 1.90 (2H, m), 2.10 - 2.20 (2H, m), 2.46 (3H, s), 2.55 (2H, m), 2.67 (4H, m), 3.01 (4H, m), 4.00 (1H, m), 6.04 (1H, d, $J = 8$ Hz), 7.24 (1H, d, $J = 8$ Hz), 7.55 (2H, m), 7.80 - 7.95 (6H, m), 8.25 (1H, s).</p>
126	7-(3,4-dihydro-3-oxo)-2H-benzoxazinyl	<p>Mass spectrum (API⁺): Found 530 (MH⁺). $C_{30}H_{35}N_5O_4$ requires 529.</p> <p>NMR ($CDCl_3$) δ: 1.10 - 1.30 (5H, m), 1.40 - 1.50 (2H, m), 1.75 - 1.85 (2H, m), 2.00 - 2.10 (2H, m), 2.46 (3H, s), 2.50 - 2.60 (2H, m), 2.64 - 2.75 (4H, m), 2.95 - 3.05 (4H, m), 3.90 (1H, m), 4.64 (2H, s), 5.79 (1H, d, $J = 8$ Hz), 6.81 (1H, d, $J = 8$ Hz), 7.20 - 7.22 (1H, m), 7.40 (2H, m), 7.72 (1H, br s), 7.83 - 7.90 (2H, m).</p>
127	5-quinolinyl(2-Me)	<p>Mass spectrum (API⁺): Found 524 (MH⁺). $C_{32}H_{37}N_5O_2$ requires 523.</p> <p>NMR ($DMSO-d_6$) δ: 1.02 - 1.10 (2H, m), 1.20 - 1.40 (5H, m), 1.75 - 1.83 (2H, m), 1.90 - 2.00 (2H, m), 2.33 (2H, m), 2.40 (3H, s), 2.55 - 2.60 (4H, m), 2.66 (3H, s), 2.90 - 3.00 (4H, m), 3.75 - 3.85 (1H, s), 7.35 - 7.37 (1H, m), 7.44 - 7.47 (1H, m), 7.57 - 7.59 (1H, m), 7.69 - 7.72 (1H, m), 7.81 - 7.85 (2H, m), 7.96 - 8.00 (1H, m), 8.41 - 8.48 (2H, m).</p>
128	-CH ₂ -(2-fluoro)phenyl	<p>Mass spectrum (API⁺): Found 491 (MH⁺). $C_{29}H_{35}FN_4O_2$ requires 490.</p>

		NMR (CDCl ₃) δ : 1.00 – 1.07 (4H, m), 1.18 – 1.23 (1H, m), 1.38 – 1.43 (2H, m), 1.72 – 1.76 (2H, m), 1.91 – 1.94 (2H, m), 2.46 (3H, s), 2.45 – 2.49 (2H, m), 2.60 – 2.64 (4H, m), 2.95 – 2.99 (4H, m), 3.54 (2H, s), 3.67 – 3.72 (1H, m), 5.25 (1H, d, J = 8 Hz), 7.04 – 7.14 (2H, m), 7.21 – 7.32 (3H, m), 7.84 – 7.86 (2H, m).
129	-CH ₂ -(2,5-difluoro)phenyl	Mass spectrum (API ⁺): Found 509 (MH ⁺). C ₂₉ H ₃₄ F ₂ N ₄ O ₂ requires 508. NMR (CDCl ₃) δ : 0.96 – 1.29 (5H, m), 1.37 – 1.46 (2H, m), 1.67 – 1.79 (2H, m), 1.93 – 1.97 (2H, m), 2.46 (3H, s), 2.44 – 2.50 (2H, m), 2.60 – 2.65 (4H, m), 2.95 – 3.05 (4H, m), 3.50 (2H, s), 3.62 – 3.76 (1H, m), 5.30 (1H, d, J = 8 Hz), 6.89 – 7.08 (2H, m), 7.21 – 7.24 (2H, m), 7.84 – 7.87 (2H, m).
130	2-indolyl	Mass spectrum (API ⁺): Found 498 (MH ⁺). C ₃₀ H ₃₅ N ₅ O ₂ requires 497. NMR (DMSO-d ₆) δ : 0.97 – 1.11 (2H, m), 1.26 – 1.50 (5H, m), 1.70 – 2.00 (4H, m), 2.39 – 2.62 (5H, m), 2.93 – 3.02 (4H, m), 3.31 – 3.40 (4H, m), 3.70 – 3.90 (1H, m), 6.98 – 7.04 (1H, m), 7.13 – 7.18 (2H, m), 7.35 – 7.43 (2H, m), 7.59 (1H, d, J = 8 Hz), 7.81 – 7.85 (2H, m), 8.20 (1H, d, J = 8 Hz), 11.50 – 11.54 (1H, s).

Table 3.



5


Example No.	R	Mass spectrum, ¹ H NMR
131	-CH ₂ -(2-benzothiophenyl)	Mass spectrum (API ⁺): Found 529 (MH ⁺). C ₃₁ H ₃₆ N ₄ O ₂ S requires 528.

		NMR (CDCl ₃) δ : 0.95 - 1.10 (4H, m), 1.18 (1H, m), 1.35 - 1.45 (2H, m), 1.74 (2H, m), 1.95 (2H, m), 2.45 (2H, m), 2.55 - 2.68 (4H, m), 2.64 (3H, s), 2.90 - 3.00 (4H, m), 3.73 (1H, m), 3.82 (2H, s), 5.46 (1H, d, J = 8 Hz), 7.16 (1H, s), 7.18 (1H, d, J = 8 Hz), 7.27 - 7.40 (2H, m), 7.72 (1H, d, J = 7 Hz), 7.76 - 7.85 (3H, m).
132	(E)-CH=CH-(3-thienyl)	Mass spectrum (API ⁺): Found 491 (MH ⁺). C ₂₈ H ₃₄ N ₄ O ₂ S requires 490. NMR (CDCl ₃) δ : 1.05 - 1.20 (4H, m), 1.24 (1H, m), 1.44 (2H, m), 1.80 (2H, m), 2.05 (2H, m), 2.50 (2H, m), 2.55 - 2.70 (7H, m), 2.90 - 3.05 (4H, m), 3.85 (1H, m), 5.35 (1H, d, J = 8 Hz), 6.18 (1H, d, J = 16 Hz), 7.19 (1H, d, J = 8 Hz), 7.21 - 7.27 (1H, m), 7.32 (1H, m), 7.43 (1H, m), 7.59 (1H, d, J = 16 Hz), 7.75 - 7.85 (2H, m).
133	5-quinolinyl	Mass spectrum (API ⁺): Found 510 (MH ⁺). C ₃₁ H ₃₅ N ₅ O ₂ requires 509. NMR (CDCl ₃) δ : 1.10 - 1.35 (5H, m), 1.48 (2H, m), 1.80 - 1.90 (2H, m), 2.10 - 2.25 (2H, m), 2.53 (2H, m), 2.65 (3H, s), 2.60 - 2.70 (4H, m), 2.99 (4H, m), 4.03 (1H, m), 5.85 (1H, d, J = 8 Hz), 7.20 (1H, d, J = 8 Hz), 7.46 (1H, dd, J = 4, 8 Hz), 7.66 (2H, m), 7.78 - 7.85 (2H, m), 8.16 (1H, d, J = 8 Hz), 8.74 (1H, d, J = 8 Hz), 8.95 (1H, m).
134	3- pyrrolo[2,3-b]pyridyl	Mass spectrum (API ⁺): Found 499 (MH ⁺). C ₂₉ H ₃₄ N ₆ O ₂ requires 498. NMR (DMSO-d ₆) δ : 0.90 - 1.10 (2H, m), 1.20 - 1.50 (5H, m), 1.70 - 1.90 (4H, m), 2.40 - 2.60 (6H, m), 2.65 (3H, s), 2.93 (4H, m), 3.75 (1H, m), 7.14 (1H, dd, J = 4, 8 Hz), 7.29 (1H, d, J = 8 Hz), 7.60 - 7.80 (3H, m), 8.14 (1H, s), 8.23 (1H, m), 8.43 (1H, m), 11.99 (1H, s).
135	8-(1,4-dihydro-4-oxo)quinolyl	Mass spectrum (API ⁺): Found 526 (MH ⁺). C ₃₁ H ₃₅ N ₅ O ₃ requires 525.

		NMR (CDCl ₃) δ : 1.10 - 1.20 (2H, m), 1.20 - 1.34 (3H, m), 1.42 - 1.50 (2H, m), 1.80 - 1.90 (2H, m), 2.05 - 2.15 (2H, m), 2.50 (2H, m), 2.65 (3H, s), 2.65 - 2.70 (4H, m), 2.98 (4H, m), 3.95 (1H, m), 6.30 (1H, d, J = 8 Hz), 6.33 (1H, dd, J = 2, 8 Hz), 7.20 (1H, d, J = 8 Hz), 7.31 (1H, t, J = 8 Hz), 7.67 (1H, t, J = 8 Hz), 7.81 (3H, m), 8.55 (1H, d, J = 8 Hz), 12.20 (1H, br s).
136	3-(3-(5-methyl)-1,2,4-oxadiazolyl)phenyl	Mass spectrum (API ⁺): Found 541 (MH ⁺). C ₃₁ H ₃₆ N ₆ O ₃ requires 540. NMR (CDCl ₃) δ : 1.10 - 1.30 (5H, m), 1.40 (2H, m), 1.83 (2H, m), 2.10 (2H, m), 2.52 (2H, m), 2.60 - 2.70 (10H, m), 2.98 (4H, m), 3.96 (1H, m), 6.00 (1H, d, J = 8 Hz), 7.20 (1H, d, J = 8 Hz), 7.57 (1H, t, J = 8 Hz), 7.75 - 7.82 (2H, m), 7.97 (1H, d, J = 8 Hz), 8.17 (1H, d, J = 8 Hz), 8.32 (1H, s).
137	(E)-CH=CH(4-fluoro)phenyl	Mass spectrum (API ⁺): Found 503 (MH ⁺). C ₃₀ H ₃₅ FN ₄ O ₂ requires 502. NMR (CDCl ₃) δ : 1.10 - 1.80 (4H, m), 1.25 (1H, m), 1.44 (2H, m), 1.78 (2H, m), 2.06 (2H, m), 2.50 (2H, m), 2.60 - 2.70 (7H, m), 2.90 - 3.00 (4H, m), 3.85 (1H, m), 5.39 (1H, d, J = 8 Hz), 6.26 (1H, d, J = 16 Hz), 7.05 (2H, t, J = 8 Hz), 7.20 (1H, d, J = 8 Hz), 7.47 (2H, m), 7.57 (1H, d, J = 16 Hz), 7.80 - 7.90 (2H, m).
138	(E)-CH=CH-(3-F)phenyl	Mass spectrum (API ⁺): Found 503 (MH ⁺). C ₃₀ H ₃₅ FN ₄ O ₂ requires 502. NMR (CDCl ₃) δ : 1.05 - 1.20 (4H, m), 1.20 - 1.30 (1H, m), 1.40 - 1.50 (2H, m), 1.75 - 1.85 (2H, m), 2.00 - 2.10 (2H, m), 2.45 - 2.55 (2H, m), 2.60 - 2.70 (7H, m), 2.90 - 3.05 (4H, m), 3.80 - 3.90 (1H, m), 5.41 (1H, d, J = 8 Hz), 6.33 (1H, d, J = 15 Hz), 6.95 - 7.05 (1H, m), 7.13 - 7.20 (2H, m), 7.20 - 7.25 (1H, m), 7.27 - 7.35 (1H, m), 7.56 (1H, d, J = 15 Hz), 7.75 - 7.85 (2H, m).
139	(E)-CH=CH-(2-	Mass spectrum (API ⁺): Found 503 (MH ⁺).

	F)phenyl	<p>$C_{30}H_{35}FN_4O_2$ requires 502.</p> <p>NMR ($CDCl_3$) δ: 1.06 - 1.30 (5H, m), 1.40 - 1.50 (2H, m), 1.75 - 1.85 (2H, m), 2.00 - 2.10 (2H, m), 2.45 - 2.55 (2H, m), 2.60 - 2.70 (7H, m), 2.90 - 3.00 (4H, m), 3.80 - 3.90 (1H, m), 5.42 (1H, d, $J = 8$ Hz), 4.49 (1H, d, $J = 15$ Hz), 7.10 - 7.22 (3H, m), 7.26 - 7.31 (1H, m), 7.40 - 7.50 (1H, m), 7.66 (1H, d, $J = 15$ Hz), 7.75 - 7.85 (2H, m).</p>
--	----------	--

Table 4.




Example	R ¹	A	Mass spectrum
140	Me	<i>trans</i> -CH=CHC ₆ H ₄ (2-F)	Found: 499 (MH ⁺) $C_{28}H_{35}N_2SO_3F$ requires 498
141	Me	-C ₆ H ₄ (3-(2-(4-Methyl)-oxazolyl))	Found: 536 (MH ⁺) $C_{30}H_{37}N_3SO_4$ requires 535
142	Me	-C ₆ H ₄ (3-trifluoromethyl)	Found: 523 (MH ⁺) $C_{27}H_{33}N_2SO_3F_3$ requires 522
143	Me	5-quinolinyl(8-Cl, 2-Me)	Found: 554 (MH ⁺) $C_{30}H_{36}N_3SO_3Cl$ requires 553
163	Me	5-quinolinyl(8-F, 2-Me)	Found: 538 (MH ⁺) $C_{30}H_{36}FN_3O_3S$ requires 537

The substituted benzazepines required as intermediates for the compounds of Table 5 were prepared from the compounds of Descriptions 8, 9, or 13, using standard methods for functional group transformation and heterocyclic ring synthesis or by palladium-catalysed cross-coupling reactions.

5

Table 5



Example	R¹	A	Mass Spectrum (API⁺)
144		-CH₂Ph(2-F)	Found 490 (MH⁺). C₃₀H₃₆FN₃O₂ requires 489.
145		Ph(4-F)	Found 502 (MH⁺). C₃₁H₃₆FN₃O₂ requires 501.
146		Ph(4-F)	Found 502 (MH⁺). C₃₁H₃₆FN₃O₂ requires 501.
147		-CH₂Ph(4-F)	Found 490 (MH⁺). C₃₀H₃₆FN₃O₂ requires 489.
148	2-pyridyl	Ph(4-F)	Found 498 (MH⁺). C₃₂H₃₆FN₃O requires 497.
149	2-pyrimidinyl	Ph(4-F)	Found 499 (MH⁺). C₃₁H₃₅FN₄O requires 498.
151		Ph(4-F)	Found 518 (MH⁺). C₃₂H₄₀FN₃O₂ requires 517.
152		3-(7-aza)indolyl	Found 514 (MH⁺). C₃₁H₃₉N₅O₂ requires 513.
153	5-pyrimidinyl	3-(3-(5-methyl)-1,2,4-oxadiazolyl)phenyl	Found 537 (MH⁺). C₃₂H₃₆N₆O₂ requires 536.
154	5-pyrimidinyl	5-quinolinyl(2-Me)	Found 520 (MH⁺). C₃₃H₃₇N₅O requires 519.
155	5-pyrimidinyl	Ph(4-F)	Found 499 (MH⁺). C₃₁H₃₅FN₄O requires 498.
156		5-quinolinyl(2-Me)	Found 523 (MH⁺). C₃₃H₃₈N₄O₂ requires 522.
157		<i>trans</i> -CH=CHC₆H₄(2-CN)	Found 509 (MH⁺). C₃₂H₃₆N₄O₂ requires 508.
158		<i>trans</i> -CH=CHC₆H₄(3-CN)	Found 509 (MH⁺). C₃₂H₃₆N₄O₂ requires 508.
159		<i>trans</i> -CH=CHC₆H₄(4-CN)	Found 509 (MH⁺). C₃₂H₃₆N₄O₂ requires 508.

161	MeSO ₂ O-	5-quinoliny(8-F, 2-Me)	Found 554 (MH ⁺). C ₃₀ H ₃₆ FN ₃ O ₄ S requires 553.
162	MeSO ₂ O-	-C ₆ H ₄ (3-(2-(5-Methyl)-oxazolyl))	Found 552 (MH ⁺). C ₃₀ H ₃₇ N ₃ O ₅ S requires 551.

Example 12

5 ***trans*-(E)-7-Cyano-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine**

10 A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-cyano-2,3,4,5-tetrahydro-1H-3-benzazepine (103 mg, 0.35 mmol), 4-fluorocinnamic acid (58 mg, 0.35 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (67 mg, 0.35 mmol), and 1-hydroxybenzotriazole (20 mg, 0.15 mmol) in dichloromethane (8 ml) was shaken at room temperature for 16 h. The reaction mixture was washed with saturated sodium bicarbonate (4 ml). The resulting precipitate was collected by filtration, washed with water (2 x 10 ml), and dried to give the title compound (87 mg, 56%) as a colourless solid.

15 Mass spectrum (API⁺): Found 446 (MH⁺). C₂₈H₃₂FN₃O requires 445.

¹H NMR (DMSO-d₆) δ: 0.94 – 1.31 (8H, m), 1.81 (4H, m), 2.40 (5H, m), 3.04 (4H, m), 3.63 (1H, m), 6.54 (1H, d, J = 16 Hz), 7.32 (4H, m), 7.59 (4H, m), 7.99 (1H, d, J = 8 Hz).

Example 13

25 ***trans*-7-Cyano-3-(2-(1-(4-(3-pyrrolo[2,3-b]pyridyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine**

30 A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-cyano-2,3,4,5-tetrahydro-1H-3-benzazepine (103 mg, 0.35 mmol), 3-pyrrolo[2,3-b]pyridyl carboxylic acid (56 mg, 0.35 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (67 mg, 0.35 mmol) and 1-hydroxybenzotriazole (20 mg, 0.15 mmol) in dichloromethane (8 ml) was shaken at room temperature for 16 h. The reaction mixture was washed with saturated aqueous sodium bicarbonate (4 ml). The resulting precipitate was collected by filtration, washed with water (2 x 10 ml) and dried to give the title compound (81 mg, 0.18 mmol, 53%) as a colourless solid.

35 Mass spectrum (API⁺): Found 442 (MH⁺). C₂₇H₃₁N₅O requires 441.

¹H NMR (DMSO-d₆) δ: 1.02 (2H, m), 1.15 – 1.45 (6H, m), 1.81 (4H, m), 2.50 (5H, m), 2.91 (4H, m), 3.73 (1H, m), 7.14 (1H, m), 7.32 (1H, d, J = 8 Hz), 7.57 (2H, m), 7.73 (1H, d, J = 8 Hz), 8.16 (1H, m), 8.25 (1H, m), 8.42 (1H, m), 12.03 (1H, br s).

Example 14***trans*-7-Cyano-3-(2-(1-(4-(3-(3-(5-methyl)-1,2,4-oxadiazolyl)benzoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

5 A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-cyano-2,3,4,5-tetrahydro-1*H*-3-benzazepine (103 mg, 0.35 mmol), 3-(3-(5-methyl)-1,2,4-oxodiazolyl)benzoic acid (71 mg, 0.35 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (67 mg, 0.35 mmol) and 1-hydroxybenzotriazole (20 mg, 0.15 mmol) in dichloromethane (8 ml)
10 was shaken at room temperature for 16 h. The reaction mixture was washed with saturated aqueous sodium bicarbonate (4 ml). The organic layer was pipetted onto a 10 g pre-packed silica column and eluted with 30 – 100% ethyl acetate in hexane. The fractions containing the title compound were combined and evaporated *in vacuo* to give the title compound (119 mg, 71%) as a colourless solid.

15 Mass spectrum (API⁺): Found 484. C₂₉H₃₃N₅O₂ requires 483.

¹H NMR (CDCl₃) δ: 1.08 – 1.35 (5H, m), 1.45 (2H, m), 1.84 (2H, m), 2.12 (2H, m), 2.50 (2H, m), 2.62 (4H, m), 2.68 (3H, s), 2.96 (4H, m), 3.95 (1H, m), 6.02 (1H, d, J = 8 Hz),
20 7.17 (1H, d, J = 8 Hz), 7.41 (2H, m), 7.57 (1H, t, J = 8 Hz), 7.98 (1H, m), 8.17 (1H, m), 8.32 (1H, m).

Example 15***trans*-(*E*)-7-Cyano-3-(2-(1-(4-(5-quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-benzazepine**

A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-cyano-2,3,4,5-tetrahydro-1*H*-benzazepine (0.10 g, 0.34 mmol), quinoline-5-carboxylic acid (0.057 g, 0.37 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (0.065 g, 0.34 mmol), 1-
30 hydroxybenzotriazole (catalytic amount) and dichloromethane (8 ml) was shaken for 16 h. Saturated sodium bicarbonate (4 ml) was then added and the mixture shaken for 0.25 h. Chromatography on the organic layer on silica eluting with a gradient of 30 - 100% ethyl acetate in hexane and then 0 - 10% methanol in ethyl acetate gave the title compound (0.130 g, 86%).

35 Mass spectrum (API⁺) Found 453 (MH⁺). C₂₉H₃₂N₄O requires 452.

¹H NMR (CDCl₃) δ: 1.12 - 1.35 (5H, m), 1.41 - 1.51 (2H, m), 1.83 - 1.89 (2H, m), 2.15 - 2.24 (2H, m), 2.48 - 2.55 (2H, m), 2.60 - 2.66 (4H, m), 2.91 - 2.99 (4H, m), 3.97 - 4.13

(1H, m), 5.86 (1H, d, J = 8 Hz), 7.18 (1H, d, J = 8 Hz), 7.37 - 7.49 (3H, m), 7.63 - 7.70 (2H, m), 8.15 - 8.20 (1H, m), 8.71 - 8.76 (1H, m), 8.94 - 8.96 (1H, m).

Example 16

5

***trans*-(E)-7-Cyano-3-(2-(1-(4-(3-(3-acetylamino)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine**

A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-cyano-2,3,4,5-tetrahydro-1H-3-benzazepine (0.10 g, 0.34 mmol), 3-acetamido cinnamic acid (0.076 g, 0.42 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (0.071 g, 0.42 mmol), 1-hydroxybenzotriazole (catalytic amount) and dichloromethane (8 ml) was shaken for 16 h. Saturated sodium bicarbonate (4 ml) was then added and the mixture shaken for 0.25 h. The precipitated solid was filtered off and washed with water then diethyl ether, and dried to give the title compound (0.12 g, 74%) as a colourless solid.

Mass spectrum (API⁺): Found 485 (MH⁺). C₃₀H₃₆N₄O₂ requires 484.

¹H NMR (CDCl₃ + CD₃OD) δ: 1.02 - 1.35 (5H, m), 1.35 - 1.50 (2H, m), 1.77 - 1.82 (2H, m), 2.00 - 2.04 (2H, m), 2.17 (3H, s), 2.47 - 2.55 (6H, m), 2.93 - 2.99 (4H, m), 3.70 - 3.85 (1H, m), 6.41 (1H, d, J = 15 Hz), 7.17 - 7.30 (4H, m), 7.38 - 7.43 (3H, m), 7.50 (1H, d, J = 16 Hz), 7.80 (1H, s).

Example 17

25

***trans*-7-Cyano-3-(2-(1-(4-(6-(3,4-dihydro-3-oxo)-2H-benzoxazinyl)carboxamido)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-benzazepine**

A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-cyano-2,3,4,5-tetrahydro-1H-3-benzazepine (0.10 g, 0.34 mmol), 2,3-dihydro-3-oxo-4H-benzoxazine-6-carboxylic acid (0.072 g, 0.42 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (0.071 g, 0.42 mmol), 1-hydroxybenzotriazole (catalytic amount) and dichloromethane (8 ml) was shaken for 16 h. Saturated sodium bicarbonate (4 ml) was then added and the mixture shaken for 0.25 h. The precipitated solid was filtered off and washed with water then diethyl ether, and dried to give the title compound (0.16 g, 100%) as a colourless solid.

Mass spectrum (API⁺): Found 473 (MH⁺). C₂₈H₃₂N₄O₃ requires 472.

¹H NMR (DMSO-d₆) δ: 0.95 - 1.50 (7H, m), 1.75 - 1.95 (4H, m), 2.40 - 2.65 (6H, m),
 5 2.93 - 3.05 (4H, m), 3.69 - 3.82 (1H, m), 4.67 (2H, s), 7.02 (1H, d, J = 8 Hz), 7.39 (1H, d,
 J = 8 Hz), 7.46 - 7.50 (2H, m), 7.65 (2H, m), 8.13 (1H, d, J = 8 Hz).

Example 18

10 ***trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(6-(1,2-dihydro-2-oxo)quinolinyl)propenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-cyano-2,3,4,5-tetrahydro-1*H*-3-benzazepine (288 mg, 0.97 mmol), *trans*-3-(6-(1,2-dihydro-2-oxo)quinolinyl)-propenoic acid (250 mg, 1.16 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide
 15 hydrochloride (204 mg, 1.07 mmol), 1-hydroxybenzotriazole (catalytic amount) and DMF (20 ml) was shaken for 18 h. Saturated sodium bicarbonate (8 ml) was then added and the mixture shaken for 0.25 h. The resulting precipitate was filtered and dried *in vacuo* to give the title compound (370 mg, 77%) as a colourless solid.

20 Found: 495 (MH⁺). C₃₁H₃₄N₄O₂ requires 494.

¹H NMR (DMSO-d₆) δ: 0.94 - 1.05 (2H, m), 1.10 - 1.30 (3H, m), 1.30 - 1.40 (2H, m),
 1.74 - 1.80 (2H, m), 1.80 - 1.88 (2H, m), 2.44 (2H, t, J = 7.5 Hz), 2.45 - 2.55 (4H, m),
 2.85 - 2.95 (4H, m), 3.55 - 3.65 (1H, m), 6.50 - 6.60 (2H, m), 7.28 - 7.35 (2H, m), 7.40
 25 (1H, d, J = 16 Hz), 7.55 - 7.60 (2H, m), 7.68 - 7.72 (1H, m), 7.81 (1H, s), 7.93 (1H, d, J =
 16 Hz), 7.94 - 8.00 (2H, m).

Example 19

30 ***trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-fluoro-4-acetyl amino)phenylpropenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-cyano-2,3,4,5-tetrahydro-1*H*-3-benzazepine (150 mg, 0.51 mmol), (*E*)-(2-fluoro-4-acetyl amino)phenylpropenoic acid (113 mg, 0.51 mmol), EDC. hydrochloride (97 mg, 0.51 mmol) and 1-

hydroxybenzotriazole in dichloromethane (10 ml) was shaken at room temperature for 16 h. The reaction mixture was washed with saturated aqueous sodium bicarbonate (4 ml) and the precipitate collected by filtration and then re-suspended in water and filtered before drying *in vacuo* to give the title compound as an off white solid (200 mg, 79%).

5

Mass spectrum (API⁺): Found 503. C₃₀H₃₅N₄O₂ requires 502.

¹H NMR δ (DMSO-d₆ + TFA): 0.95 - 1.34 (5H, m), 1.61 (2H, m), 1.82 (4H, m), 2.07 (3H, s), 3.06 (2H, m), 3.18 (6H, m), 3.68 (3H, m), 6.59 (1H, d, J = 16 Hz), 7.34 (2H, m),
10 7.39 - 7.63 (3H, m), 7.72 (2H, m), 8.03 (1H, d, J = 8 Hz), 9.74 (1H, br s), 10.29 (1H, s).

Example 20

15 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(8-(1,2-dihydro-2-oxo)quinolinyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine

A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-cyano-2,3,4,5-tetrahydro-1*H*-3-benzazepine (0.25 g, 0.84 mmol), (*E*)-3-(8-(1,2-dihydro-2-oxo)quinolinyl)propenoic acid (0.27 g, 1.2 mmol), EDC. hydrochloride (0.3 g, 1.5 mmol) and 1-hydroxybenzotriazole (50 mg) in DMF (10 ml) was allowed to stir at 80 °C for 4 h, then poured into water (500
20 ml). The precipitate was collected by filtration and then re-suspended in aqueous sodium bicarbonate solution. Resulting solid was collected by filtration, then washed with water and diethyl ether, then was dried *in vacuo* to give the title compound (0.42 g, 95 %) as a yellow solid.

25 Found: 495 (MH⁺). C₃₁H₃₄N₄O₂ requires 494.

δ (DMSO-d₆ + TFA): 1.00 - 1.15 (2H, m), 1.15 - 1.30 (3H, m), 1.50 - 1.70 (2H, m), 1.70 - 1.85 (2H, m), 1.85 - 1.95 (2H, m), 2.95 - 3.30 (8H, m), 3.60 - 3.80 (3H, m), 6.45 - 6.60 (2H, m), 7.23 (1H, t, J = 8 Hz), 7.46 (1H, d, J = 8 Hz), 7.60 - 7.80 (4H, m), 7.94 (1H, d, J
30 = 10 Hz), 7.95 - 8.10 (3H, m), 9.70 (1H, br s).

Example 21

***trans*-7-Cyano-3-(2-(1-(4-(5-(8-fluoro)quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine**

A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-cyano-2,3,4,5-tetrahydro-1H-3-benzazepine (0.162 g, 0.545 mmol), 8-fluoroquinoline-5-carboxylic acid (0.115 g, 0.6 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.115 g, 0.6 mmol) and 1-hydroxybenzotriazole hydrate (0.01 g, 0.065 mmol) in dichloromethane (7 ml) was shaken for 18 h. Saturated aqueous sodium hydrogen carbonate (6 ml) was added and shaking continued for 0.5 h. The organic layer was separated and pipetted onto a column of silica (10 g). Elution with 30 - 100% ethyl acetate - hexane gradient then 1 - 10% methanol - ethyl acetate gradient yielded the title compound as a colourless solid (0.22 g, 85%).

Mass spectrum (API⁺): Found 471 (MH⁺). C₂₉H₃₁FN₄O requires 470.

¹H NMR (CDCl₃) δ: 1.05 - 1.40 (5H, m), 1.45 (2H, m), 1.85 (2H, m), 2.20 (2H, m), 2.55 (2H, m), 2.63 (4H, m), 2.96 (4H, m), 4.00 (1H, m), 5.86 (1H, d, J = 8 Hz), 7.17 (1H, d, J = 8 Hz), 7.30 - 7.45 (3H, m), 7.54 (1H, m), 7.62 (1H, m), 8.80 (1H, d, J = 8 Hz), 9.01 (1H, m).

Example 22

***trans*-7-Acetyl-3-(2-(1-(4-(5-quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine**

A mixture of 7-acetyl-*trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine (0.105 g, 0.334 mmol), quinoline-5-carboxylic acid (0.064 g, 0.368 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.071 g, 0.368 mmol) and 1-hydroxybenzotriazole hydrate (0.01 g, 0.065 mmol) in dichloromethane (6 ml) was shaken for 18 h. Saturated aqueous sodium hydrogen carbonate (6 ml) was added and shaking continued for a further 0.5 h. The organic layer was separated and pipetted onto a column of silica (10 g). Elution with 30 - 100% ethyl acetate - hexane gradient then 1 - 10% methanol - ethyl acetate gradient gave the title compound as a colourless solid (0.1 g, 64%).


Mass spectrum (API⁺): Found 470 (MH⁺); C₃₀H₃₅N₃O₂ requires 469.

¹H NMR (CDCl₃) δ: 1.10 - 1.40 (5H, m), 1.48 (2H, m), 1.86 (2H, m), 2.33 (2H, m), 2.55 (2H, m), 2.58 (3H, s), 2.65 (4H, m), 2.98 (4H, m), 4.02 (1H, m), 5.88 (1H, d, J = 8 Hz), 7.17 (1H, d, J = 8 Hz), 7.20 (1H, m), 7.55 - 7.75 (4H, m), 8.15 (1H, m), 8.75 (1H, d, J = 8 Hz), 8.95 (1H, m).

5

Example 23

***trans*-3-(2-(1-(4-(5-(2-Methyl)quinolinyl)carboxamido)cyclohexyl)ethyl)-7-methylsulphonyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



10 A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-methylsulphonyl-2,3,4,5-tetrahydro-1H-3-benzazepine (100 mg, 0.29 mmol), 2-methyl-quinoline-5-carboxylic acid (64 mg, 0.34 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (59 mg, 0.31 mmol) and 1-hydroxybenzotriazole (cat. amt.) in dichloromethane (10 ml) was shaken at room temperature for 18 h. A saturated solution of sodium bicarbonate (4 ml)
15 was then added and the mixture shaken for 0.25 h. The organic layer was then applied directly to a silica column eluted with a gradient of 30 - 100% ethyl acetate in hexane and then 0 - 10% methanol in ethyl acetate to give the title compound (95 mg, 66 %) as a colourless solid.

20 ¹H NMR δ (CDCl₃) 1.15 - 1.30 (5H, m), 1.44 - 1.50 (2H, m), 1.82 - 1.88 (2H, m), 2.15 - 2.20 (2H, m), 2.53 (2H, t, J = 7.6 Hz), 2.62 - 2.68 (4H, m), 2.75 (3H, s), 2.98 - 3.02 (4H, m), 3.04 (3H, s), 3.95 - 4.05 (1H, m), 5.84 (1H, d, J = 8.2 Hz), 7.28 (1H, d, J = 7.9 Hz), 7.35 (1H, d, J = 8.8 Hz), 7.56 - 7.70 (4H, m), 8.08 (1H, d), 8.62 (1H, d).

Mass spectrum: API⁺ 520 (MH⁺): C₃₀H₃₇N₃SO₃ requires 519.

25

Example 24

***trans*-3-(2-(1-(4-(3-(3-(5-Methyl)-1,2,4-oxadiazolyl)benzoyl)amino)cyclohexyl)ethyl)-7-methylsulphonyl-2,3,4,5-tetrahydro-1H-3-benzazepine**

A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)-7-methylsulphonyl-2,3,4,5-tetrahydro-1H-3-benzazepine (100 mg, 0.29 mmol), 3-(3-(5-methyl)-1,2,4-oxadiazolyl)-benzoic acid (69 mg, 0.34 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (59 mg, 0.31 mmol) and 1-hydroxybenzotriazole (cat. amt.) in dichloromethane (10 ml) was
30

shaken at room temperature for 18 h. A saturated solution of sodium bicarbonate (4 ml) was then added and the mixture shaken for 0.25 h. The organic layer was then applied directly to a silica column eluted with a gradient of 30 - 100% ethyl acetate in hexane and then 0 - 10% methanol in ethyl acetate to give the title compound (103 mg, 69 %) as a colourless solid.

^1H NMR δ (CDCl_3): 1.08 - 1.30 (5H, m), 1.40 - 1.46 (2H, m), 1.80 - 1.85 (2H, m), 2.08 - 2.15 (2H, m), 2.52 (2H, t, $J = 7.8$), 2.60 - 2.65 (4H, m), 2.68 (3H, s), 2.98 - 3.02 (4H, m), 3.05 (3H, s), 3.90 - 4.00 (1H, m), 6.01 (1H, d, $J = 8.0$ Hz), 7.28 (1H, d, $J = 7.28$ Hz), 7.57 (1H, t, $J = 7.8$ Hz), 7.65 - 7.70 (2H, m), 8.0 (1H, d), 8.19 (1H, d, $J = 7.7$ Hz), 8.32 (1H, s).

Mass spectrum: API^+ 537 (MH^+): $\text{C}_{29}\text{H}_{36}\text{N}_4\text{SO}_4$ requires 536.

Example 25

***trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-pyrrolo[2,3-b]pyridyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine, hydrochloride**

A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)-7-(5-(3-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1H-3-benzazepine (21.0 g, 59.3 mmol), pyrrolo[2,3-b]pyridyl-3-carboxylic acid (10.57 g, 65.2 mmol), EDC hydrochloride (12.46 g, 64.4 mmol) and HOBT (0.5 g) in CH_2Cl_2 (630 ml) and DMF (84 ml) was stirred at ambient temperature for 16 h. Saturated aqueous sodium bicarbonate (350 ml) was added and the mixture stirred for 0.25 h. The precipitate was collected by filtration, washed in turn with water and diethyl ether and dried *in vacuo* to give the free base of the title compound (18.0 g, 61 %).


Mass spectrum (API^+): Found 499 (MH^+). $\text{C}_{29}\text{H}_{34}\text{N}_6\text{O}_2$ requires 498.

NMR ($\text{DMSO}-d_6$) δ : 0.90 - 1.10 (2H, m), 1.10 - 1.40 (5H, m), 1.70 - 1.90 (4H, m), 2.40 - 2.70 (6H, m), 2.96 (3H, s), 3.31 (4H, m), 3.89 (1H, m), 7.15 (1H, m), 7.36 (1H, d, $J = 8$ Hz), 7.71 (1H, d, $J = 8$ Hz), 7.75 - 7.85 (2H, m), 8.12 (1H, s), 8.20 (1H, s), 8.35 (1H, d, $J = 8$ Hz), 12.02 (1H, br s).

To a suspension of the above free base (18.0 g, 36 mmol) in 10% methanol-dichloromethane (500 ml) was added a 1M solution of HCl in diethyl ether (37.08 ml). The resulting solution was evaporated *in vacuo* and the residue crystallised from methanol to give the title compound as a colourless solid (12.5 g, m.p. 275 - 276 °C).

Example 26

***trans*-(*E*)-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(4-fluoro)phenyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine**



- 5 *Alternative Name*: (2*E*)-3-(4-fluorophenyl)-*N*-[*trans*-4-[2-[2,3,4,5-tetrahydro-7-(5-methyl-1,2,4-oxadiazol-3-yl)-1*H*-3-benzazepin-3-yl]ethyl]cyclohexyl]-2-propenamide

A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-(3-(5-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine (0.1 g, 0.28 mmol), (*E*)-4-fluorocinnamic acid (0.046 g, 0.28 mmol), EDC hydrochloride (0.06 g, 0.31 mmol) and HOBT (0.015 g) in dichloromethane (8 ml) was stirred at ambient temperature for 64 h, then was washed with saturated aqueous sodium bicarbonate (4 ml). The organic phase was purified by silica gel chromatography eluting with 0 - 10 % methanol in ethyl acetate, to give the title compound (0.12 g, 85 %) as a colourless solid.

15

Mass spectrum (AP⁺): Found 503 (MH⁺). C₃₀H₃₅FN₄O₂ requires 502.

NMR (CDCl₃) δ: 1.10 - 1.80 (4H, m), 1.25 (1H, m), 1.44 (2H, m), 1.78 (2H, m), 2.06 (2H, m), 2.50 (2H, m), 2.60 - 2.70 (7H, m), 2.90 - 3.00 (4H, m), 3.85 (1H, m), 5.39 (1H, d, J = 8 Hz), 6.26 (1H, d, J = 16 Hz), 7.05 (2H, t, J = 8 Hz), 7.20 (1H, d, J = 8 Hz), 7.47 (2H, m), 7.57 (1H, d, J = 16 Hz), 7.80 - 7.90 (2H, m).

A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-(3-(5-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine (16.0g, 0.045 mol), (*E*)-4-fluorocinnamic acid (7.5g, 0.045 mol), EDC hydrochloride (9.53 g, 0.050 mol), and HOBT (0.78g, 0.006 mol) in dichloromethane (0.78L) was stirred under argon at ambient temperature for 11 h. Saturated aqueous sodium bicarbonate (1L) was added and after stirring for 0.25h, the mixture was filtered and the solid washed with saturated aqueous sodium bicarbonate (2 x 0.25L), water (3 x 0.25L), diethyl ether (3 x 0.25L) and dried *in vacuo* to afford the title compound (18.4g, 81%) as a colourless solid.

The filtrate was separated and the aqueous layer extracted with dichloromethane (2 x 0.3L). The combined extracts were dried and evaporated *in vacuo* to afford a pale yellow

solid (4.5g). Sequential trituration with dichloromethane (0.08L), saturated aqueous sodium bicarbonate (1 x 0.5L; 2 x 0.2L), water (3 x 0.2L), and diethyl ether (3 x 0.2L) followed by drying *in vacuo* afforded the title compound (2.8g, 12%) as a colourless solid.

5

Both batches had spectroscopic data identical to that described above.

To a solution of the free base obtained above (21.2g, 0.042 mol) in dichloromethane (0.55L) and methanol (0.1L) was added 1M hydrogen chloride in diethyl ether (0.051L, 10 0.05mol). The resulting solution was evaporated *in vacuo* and the residue crystallised from methanol to afford (2*E*)-3-(4-fluorophenyl)-*N*-[*trans*-4-[2-[2,3,4,5-tetrahydro-7-(5-methyl-1,2,4-oxadiazol-3-yl)-1*H*-3-benzazepin-3-yl]ethyl]cyclohexyl]-2-propenamide monohydrochloride (19.8g, 91%) as a colourless solid m.p. 259-261°C.

15 NMR (D₆-DMSO) δ: 1.00 - 1.09 (2H, m), 1.15 - 1.28 (3H, m), 1.60 - 1.70 (2H, m), 1.70 - 1.80 (2H, m), 1.80 - 1.90 (2H, m), 2.66 (3H, s), 2.95 - 3.25 (6H, m), 3.35 - 3.50 (2H, m), 3.55 - 3.75 (3H, m), 6.55 (1H, d, J = 16 Hz), 7.22 - 7.27 (2H, m), 7.39 (1H, d, J = 16Hz), 7.40 - 7.45 (1H, m), 7.55 - 7.64 (2H, m), 7.80 - 7.85 (1H, m), 7.87 (1H, s), 7.95 - 8.05 (1H, m), 10.60 (1H, br s).

20

Example 27

***trans*-(*E*)-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(4-fluoro)phenyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine**

A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-(5-(3-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine (0.1 g, 0.28 mmol), (*E*)-4-fluorocinnamic acid (0.046 g, 0.28 mmol) EDC hydrochloride (0.06 g, 0.31 mmol) and HOBT (0.015 g) in dichloromethane (8 ml) was stirred at ambient temperature for 64 h, then was washed with saturated aqueous sodium bicarbonate (4 ml). The organic phase was purified by silica gel chromatography eluting with 0 - 10 % methanol in ethyl acetate to give the title compound (0.12 g, 85 %) as a colourless solid.

30

Mass spectrum (API⁺): Found 503 (MH⁺). C₃₀H₃₅FN₄O₂ requires 502.

NMR (CDCl₃) δ: 1.10 - 1.30 (5H, m), 1.40 - 1.47 (2H, m), 1.78 - 1.82 (2H, m), 2.00 - 2.10 (2H, m), 2.46 (3H, s), 2.47 - 2.52 (2H, m), 2.60 - 2.70 (4H, m), 2.95 - 3.05 (4H, m),

35

3.86 (1H, m), 5.38 (1H, d, J = 8 Hz), 6.26 (1H, d, J = 16 Hz), 7.05 (2H, t, J = 8 Hz), 7.24 (1H, d, J = 8 Hz), 7.47 (2H, dd, J = 5, 8 Hz), 7.57 (1H, d, J = 16 Hz), 7.80 - 7.90 (2H, m).

Example 28

5 ***trans*-(E)-7-(5-(3-Methyl)isoxazolyl)-3-(2-(1-(4-(3-(4-fluoro)phenyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine**

- A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-(5-(3-methyl)isoxazolyl)-2,3,4,5-tetrahydro-1H-3-benzazepine (0.1 g, 0.28 mmol), 4-fluorophenylacetic acid (0.044 g, 0.28 mmol), EDC hydrochloride (0.065 g, 0.31 mmol) and HOBT (0.02 g) in
- 10 CH₂Cl₂ (8 ml) was stirred at ambient temperature for 16 h, then was washed with saturated sodium bicarbonate (4 ml). The organic phase was purified by silica gel chromatography eluting with 0 - 10 % methanol in ethyl acetate to give the title compound (0.1 g, 73 %).
- 15 Mass spectrum (API⁺): Found 490 (MH⁺). C₃₀H₃₆FN₃O₂ requires 489.
- ¹H NMR δ (CDCl₃): 0.90 - 1.10 (4H, m), 1.10 - 1.20 (1H, m), 1.30 - 1.40 (2H, m), 1.70 - 1.80 (2H, m), 1.85 - 1.95 (2H, m), 2.34 (3H, s), 2.40 - 2.50 (2H, m), 2.55 - 2.70 (4H, m), 2.90 - 3.00 (4H, m), 3.50 (2H, s), 3.65 - 3.80 (1H, m), 5.12 (1H, d, J = 8 Hz), 6.30 (1H, s),
- 20 7.03 (2H, t, J = 8 Hz), 7.15 (1H, d, J = 8 Hz), 7.19 - 7.25 (2H, m), 7.45 - 7.52 (2H, m).

Example 29

***trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(4-fluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1H-3-benzazepine**

- 25 A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-(5-(3-methyl)-1,2,4-oxadiazolyl)-2,3,4,5-tetrahydro-1H-3-benzazepine (0.1 g, 0.28 mmol), (4-fluoro)phenylacetic acid (0.044 g, 0.28 mmol), EDC hydrochloride (0.054 g, 0.28 mmol) and HOBT (0.015 g) in dichloromethane (5 ml) was shaken at ambient temperature for 16 h, and saturated aqueous sodium bicarbonate (4 ml) added. The organic phase was
- 30 purified by silica gel chromatography eluting with 30-100% ethyl acetate in hexane, then 0 - 10 % methanol in ethyl acetate gradient elution to give the title compound (0.095 g, 70 %) as a colourless solid.


Mass spectrum (API⁺): Found 491 (MH⁺). C₂₉H₃₅FN₄O₂ requires 490.

¹H NMR δ (CDCl₃): 0.90 – 1.30 (5H, m), 1.35 – 1.50 (2H, m), 1.70 – 1.80 (2H, m), 1.85 – 1.95 (2H, m), 2.46 (3H, s), 2.40 – 2.50 (2H, m), 2.55 – 2.65 (4H, m), 2.95 – 3.00 (4H, m), 3.50 (2H, s), 3.60 – 3.80 (1H, m), 5.13 (1H, d, J = 8Hz), 6.95 – 7.08 (2H, m), 7.15 – 7.30 (3H, m), 7.80 – 7.90 (2H, m).

5

Example 150

***trans*-3-(2-(1-(4-(5-(2-Methyl)quinolinyl)carboxamide)cyclohexyl)ethyl)-7-methanesulphonyloxy-2,3,4,5-tetrahydro-1H-3-benzazepine**



10

A mixture of *trans*-3-(2-(1-(4-amino)cyclohexyl)ethyl)-7-methanesulphonyloxy-2,3,4,5-tetrahydro-1H-3-benzazepine (150 mg, 0.41 mmol), 2-methyl-quinoline-5-carboxylic acid (92 mg, 0.49 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (86 mg, 0.45 mmol) and 1-hydroxybenzotriazole (cat. amt.) in dichloromethane (10 ml) was shaken at room temperature for 18 h. A saturated solution of sodium bicarbonate (4 ml) was added and the mixture shaken for 0.25 h. The organic layer was then applied directly to a silica column eluted with a gradient of 30 – 100 % ethyl acetate in hexane and then 0 - 10% methanol in ethyl acetate to give the title compound (161 mg, 74%) as a colourless solid.

20


¹H NMR (CDCl₃) δ : 1.15 - 1.30 (5H, m), 1.45 - 1.50 (2H, m), 1.82 - 1.90 (2H, m), 2.15 - 2.20 (2H, m), 2.50 - 2.55 (2H, m), 2.60 - 2.68 (4H, m), 2.75 (3H, s), 2.90 - 2.95 (4H, m), 3.13 (3H, s), 3.95 - 4.05 (1H, m), 5.82 (1H, d, J = 8.2 Hz), 7.00 - 7.03 (2H, m), 7.12 (1H, d, J = 7.8 Hz), 7.35 (1H, d, J = 8.8 Hz), 7.55 - 7.70 (2H, m), 8.08 (1H, d, J = 8.3 Hz), 8.61 (1H, d).

25

Mass Spectrum (AP⁺) : Found 536 (MH⁺). C₃₀H₃₇N₃SO₄ requires 535.

Claims :

1. A compound of formula (I) :



Formula (I)

wherein:

- R^1 represents a substituent selected from: a hydrogen or halogen atom; a hydroxy, cyano, nitro, trifluoromethyl, trifluoromethoxy, trifluoromethanesulfonyloxy, pentafluoroethyl, C_{1-4} alkyl, C_{1-4} alkoxy, aryl C_{1-4} alkoxy, C_{1-4} alkylthio, C_{1-4} alkoxy C_{1-4} alkyl, C_{3-6} cycloalkyl C_{1-4} alkoxy, C_{1-4} alkanoyl, C_{1-4} alkoxycarbonyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylsulfonyloxy, C_{1-4} alkylsulfonyl C_{1-4} alkyl, arylsulfonyl, arylsulfonyloxy, arylsulfonyl C_{1-4} alkyl, C_{1-4} alkylsulfonamido, C_{1-4} alkylamido, C_{1-4} alkylsulfonamido C_{1-4} alkyl, C_{1-4} alkylamido C_{1-4} alkyl, arylsulfonamido, arylcarboxamido, arylsulfonamido C_{1-4} alkyl, arylcarboxamido C_{1-4} alkyl, aroyl, aroyl C_{1-4} alkyl, or aryl C_{1-4} alkanoyl group; a group $R^3OCO(CH_2)_p$, $R^3CON(R^4)(CH_2)_p$, $R^3R^4NCO(CH_2)_p$ or $R^3R^4NSO_2(CH_2)_p$ where each of R^3 and R^4 independently represents a hydrogen atom or a C_{1-4} alkyl group or R^3R^4 forms part of a C_{3-6} azacycloalkane or $C_{3-6}(2-oxo)azacycloalkane$ ring and p represents zero or an integer from 1 to 4; or a group Ar^3-Z , wherein Ar^3 represents an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring and Z represents a bond, O, S, or CH_2 ;

R^2 represents a hydrogen atom or a C_{1-4} alkyl group;

q is 1 or 2;


A represents a group of the formula (a), (b) (c) or (d):




(a)



(b)



(c)



(d)

wherein

- Ar represents an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring; or an optionally substituted bicyclic ring system;

Ar^1 and Ar^2 each independently represent an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring; and

- Y represents a bond, $-NHCO-$, $-CONH-$, $-CH_2-$, or $-(CH_2)_mY^1(CH_2)_n-$, wherein Y^1 represents O, S, SO_2 , or CO and m and n each represent zero or 1 such that the sum of

m+n is zero or 1; providing that when A represents a group of formula (a), any substituent present in Ar *ortho* to the carboxamide moiety is necessarily a hydrogen or a methoxy group;

5 r and s independently represent an integer from zero to 3 such that the sum of r and s is equal to an integer from 1 to 4;

V represents a bond, O or S;
and salts thereof.

10 2. A compound according to claim 1 wherein q represents 1.

3. A compound according to any of the preceding claims wherein rings Ar, Ar¹, or Ar² are each independently optionally substituted by one or more substituents selected from: a hydrogen or halogen atom, cyano, methoxy, methylenedioxy, acetyl, acetyl amino, methylsulfonyl, methylsulfonyloxy, methylaminosulfonyl,
15 methylsulfonylamino, or methylaminocarbonyl group.

4. A compound of formula (I) which is:

20 *trans*-3-(2-(1-(4-(4-Quinoliny)l)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-3-(2-(1-(4-(3-(3-Methylsulfonyl)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-3-(2-(1-(4-(3-(4-Fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-
25 tetrahydro-1*H*-3-benzazepine;
trans-3-(2-(1-(4-(2-Indolyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-3-(2-(1-(4-(3-(3-Pyridyl)phenyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
30 *trans*-3-(2-(1-(4-Phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-3-(2-(1-(4-(3-Indolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-3-(2-(1-(4-(4-Quinoliny)l)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-
35 benzazepine;
trans-(*E*)-3-(2-(1-(4-(3-(4-Fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-6-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-6-Methoxy-3-(2-(1-(4-(4-quinoliny)l)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
40 *trans*-6-Methoxy-3-(2-(1-(4-(3-pyrrolo[2,3-*b*]pyridyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-Cyano-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

- trans*-7-Cyano-3-(2-(1-(4-(3-pyrrolo[2,3-*b*]pyridyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(3-(3-(5-methyl)-1,2,4-oxadiazolyl)benzoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 5 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(5-quinoliny)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-acetylamino)phenylpropenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 10 *trans*-7-Cyano-3-(2-(1-(4-(6-(3,4-dihydro-3-oxo)-2*H*-benzoxazinyl)carboxamido)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(6-(1,2-dihydro-2-oxo)quinoliny)propenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-fluoro-4-acetylamino)phenylpropenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 15 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(8-(1,2-dihydro-2-oxo)quinoliny)propenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(5-(8-fluoro)quinoliny)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Acetyl-3-(2-(1-(4-(5-quinoliny)carboxamido)cyclohexyl)ethyl)-2,3,4,5-
- 20 tetrahydro-1*H*-3-benzazepine;
- trans*-3-(2-(1-(4-(5-(2-Methyl)quinoliny)carboxamido)cyclohexyl)ethyl)-7-methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-3-(2-(1-(4-(3-(3-(5-Methyl)-1,2,4-oxadiazolyl)benzoyl)amino)cyclohexyl)ethyl)-7-methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 25 *trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-pyrrolo[2,3-*b*]pyridyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 30 *trans*-(*E*)-7-(5-(3-Methyl)isoxazolyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(4-fluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 35 *trans*-7-Cyano-3-(2-(1-(4-(2,5-difluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(2-naphthylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2,4-difluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-
- 40 2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2,5-difluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

- trans*-7-Cyano-3-(2-(1-(4-(3-phenylpropanoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 5 *trans*-7-Cyano-3-(2-(1-(4-(8-(1,4-dihydro-4-oxo)quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(2-naphthyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-methoxy)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 10 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-methoxy)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(4-methoxy)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 15 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-acetyl)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(4-acetyl)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-cyano)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 20 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-cyano)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(3-(5-(3-methyl)isoxazolyl)benzoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 25 *trans*-7-Cyano-3-(2-(1-(4-(7-(1,2-dihydro-2-oxo)quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*Z*)-7-Cyano-3-(2-(1-(4-(3-phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-pyridyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 30 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(1-(4-fluoro)naphthyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(6-benzodioxanyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 35 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-(5-fluoro)indolyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-(1-methyl)benzimidazolyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(7-benzofuranyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 40 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(5-(3-methyl)indolyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(6-(2,3-dihydro-2-oxo)indolyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

- trans*-7-Cyano-3-(2-(1-(2-benzofuranylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(4-(2-methyl)indolyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 5 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(5-benzimidazolyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2,3-methylenedioxy)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 10 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-(1-(2-oxo)pyrrolidinyl))phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(2-indolylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 15 *trans*-7-Cyano-3-(2-(1-(4-(2-benzothiophenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-(3-bromo)thiophenyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(3-(2-pyridyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 20 *trans*-7-Cyano-3-(2-(1-(4-(3-(5-pyrimidinyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(3-(4-cyanophenyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 25 *trans*-7-Cyano-3-(2-(1-(4-(3-(3-(5-ethyl)-1,2,4-oxadiazolyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-thiophenyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-furanyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 30 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-thiophenyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-furanyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 35 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(4-quinolinyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(5-pyrimidinyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(2,4-difluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 40 *trans*-7-Cyano-3-(2-(1-(4-(1-naphthyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Acetyl-3-(2-(1-(4-(3-pyrrolo[2,3-*b*]pyridyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

- trans*-7-Acetyl-3-(2-(1-(4-(4-fluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 5 *trans*-7-Acetyl-3-(2-(1-(4-(3-(3-(5-methyl)-1,2,4-oxadiazolyl)benzoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Acetyl-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 10 *trans*-7-Cyano-3-(2-(1-(4-(6-(2-amino)benzothiazolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(6-(2-methyl)benzothiazolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(6-(2,3-dihydro-2-oxo)indolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 15 *trans*-7-Cyano-3-(2-(1-(4-(5-(2,3-dihydro-2-oxo)indolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(4-methylaminocarbonyl)phenylpropenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(5-(2-amino)benzoxazolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 20 *trans*-7-Cyano-3-(2-(1-(4-(6-(1,2-dihydro-2-oxo)quinolinyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(7-(1,2-dihydro-2-oxo)quinolinyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Acetyl-3-(2-(1-(4-(3-(3-methoxy)phenylpropenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 25 *trans*-(*E*)-7-Acetyl-3-(2-(1-(4-(3-(2-cyano)phenylpropenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Acetyl-3-(2-(1-(4-(3-(3-thiophenyl)propenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 30 *trans*-(*E*)-7-Acetyl-3-(2-(1-(4-(3-(8-(1,2-dihydro-2-oxo)quinolinyl)propenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(3-(1-pyrazolyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(2-thiophenyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 35 *trans*-7-Cyano-3-(2-(1-(4-(3-benzothiophenyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Cyano-3-(2-(1-(4-(3-(2-(5-methyl)-1,3,4-oxadiazolyl)benzoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 40 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-naphthyl)propenoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-7-Acetyl-3-(2-(1-(4-(3-benzothiophenyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- trans*-(*E*)-7-Acetyl-3-(2-(1-(4-(3-(4-acetamido)phenylpropenoyl)amino)

- cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-Acetyl-3-(2-(1-(4-(6-(2-amino)benzothiazolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-Acetyl-3-(2-(1-(4-(8-(1,4-dihydro-4-oxo)quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
5 *trans*-(*E*)-7-Acetyl-3-(2-(1-(4-(3-(2-acetyl)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-Acetyl-3-(2-(1-(4-(2-benzothiophenyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
10 *trans*-(*E*)-7-Cyano-3-(2-(1-(4-(3-(5-(3-acetyl)indolyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-Cyano-3-(2-(1-(4-(3-(5-(3-methyl)-1,2,4-oxadiazolyl)benzoyl)amino)-cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-Cyano-3-(2-(1-(4-(5-(2-methyl)benzimidazolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
15 *trans*-7-Cyano-3-(2-(1-(4-(6-quinoxaliny)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-Cyano-3-(2-(1-(4-(3-(3-(2-acetyl)furanyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
20 *trans*-7-Cyano-3-(2-(1-(4-(6-(2-amino)benzoxazolyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-Cyano-3-(2-(1-(4-(6-(3,4-dihydro-2-oxo)-2*H*-benzoxazinyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-Cyano-3-(2-(1-(4-(3-(2-fluoro-5-acetamido)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
25 *trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(2-benzothiophenyl)acetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-thienyl)propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
30 *trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(5-quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-(5-Methyl)-1,2,4-oxadiazolyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(8-(1,4-dihydro-4-oxo)quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
35 *trans*-(*E*)-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-acetamido-2-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
40 *trans*-(*E*)-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-acetyl)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-fluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(2,4-difluoro)phenylacetamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;


- cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(2-naphthyl)acetamido)
cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
5 *trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(7-(3,4-dihydro-3-oxo)-2*H*-
benzoxazinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(5-(2-methyl)quinolinyl)
carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(2-fluoro)phenylacetamido)
cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
10 *trans*-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(2,5-difluoro)phenylacetamido)
cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(5-(3-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(2-
indolyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(2-benzothiophenyl)acetamido)
15 cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-thienyl)
propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(5-quinolinyl)
carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
20 *trans*-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-pyrrolo[2,3-*b*]pyridyl)
carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(8-(1,4-dihydro-4-oxo)quinolinyl)
carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-(5-Methyl)-1,2,4-oxadiazolyl)
25 benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(3-fluoro)phenyl
propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-(3-(5-Methyl)-1,2,4-oxadiazolyl)-3-(2-(1-(4-(3-(2-fluoro)phenyl
propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
30 *trans*-3-(2-(1-(4-(5-(2-Methyl)quinolinyl)carboxamide)cyclohexyl)ethyl)-7-
methanesulphonyloxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-7-
methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-3-(2-(1-(4-(3-(2-(4-Methyl)oxazolyl)benzoyl)amino)cyclohexyl)ethyl)-7-
35 methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-3-(2-(1-(4-(3-(trifluoromethyl)benzoyl)amino)cyclohexyl)ethyl)-7-methylsulphonyl-
2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-3-(2-(1-(4-(5-(8-Chloro-2-methyl)quinolinyl)carboxamide)cyclohexyl)ethyl)-7-
methanesulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
40 *trans*-7-(5-(3-Methyl)isoxazolyl)-3-(2-(1-(4-(2-fluoro)phenylacetamido)
cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(3-(5-Methyl)isoxazolyl)-3-(2-(1-(4-(4-fluoro)phenylacetamido)
cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-(2-(5-Methyl)oxazolyl)-3-(2-(1-(4-(3-(4-fluoro)phenyl
45 propenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

- trans*-(*E*)-7-(3-(5-Methyl)isoxazolyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-(2-pyridyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
 5 *trans*-(*E*)-7-(2-pyrimidyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-(1-Pyrrolidinylcarbonyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(1-Pyrrolidinylcarbonyl)-3-(2-(1-(4-(3-pyrrolo[2,3-*b*]pyridyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
 10 *trans*-(*E*)-7-(5-Pyrimidyl)-3-(2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(5-Pyrimidinyl)-3-(2-(1-(4-(3-(3-(5-Methyl)-1,2,4-oxadiazolyl)benzoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
 15 *trans*-7-(5-Pyrimidinyl)-3-(2-(1-(4-(5-(2-methyl)quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-7-(3-(5-Methyl)isoxazolyl)-3-(2-(1-(4-(5-(2-methyl)quinolinyl)carboxamido)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-(2-Pyridyl)-3-(2-(1-(4-(3-(2-cyano)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
 20 *trans*-(*E*)-7-(2-Pyridyl)-3-(2-(1-(4-(3-(3-cyano)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-(*E*)-7-(2-Pyridyl)-3-(2-(1-(4-(3-(4-cyano)phenylpropenoyl)amino)cyclohexyl)ethyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
 25 *trans*-3-(2-(1-(4-(5-(8-Fluoro-2-methyl)quinolinyl)carboxamido)cyclohexyl)ethyl)-7-methylsulphonyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-3-(2-(1-(4-(5-(8-Fluoro-2-methyl)quinolinyl)carboxamido)cyclohexyl)ethyl)-7-methylsulphonyloxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
trans-3-(2-(1-(4-(3-(2-(5-Methyl)oxazolyl)benzoyl)amino)cyclohexyl)ethyl)-7-methylsulphonyloxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
 30

or a salt thereof.

5. A process for preparing compounds of formula (I) which process comprises :

(a) reacting a compound of formula(II):



Formula (II)

- 40 wherein R¹, R², and q are as hereinbefore defined,

with a compound of formula (III):

A-COX


5

Formula (III)

wherein A is as hereinbefore defined and X is a halogen atom or the residue of an activated ester;

10 (b) to prepare a compound of formula (I) by reacting a compound of formula (II) with a compound A-Br, or A-I, or A-OSO₂CF₃ in the presence of carbon monoxide and a catalyst;

(c) to prepare a compound of formula (I) wherein R¹ is Ar³-Z and Z is a bond, reacting a compound of formula (IV):




15

Formula (IV)

20 wherein A, R², and q are as hereinbefore defined, one R^{1a} represents a group W wherein W is a halogen atom or a trifluoromethylsulfonyloxy group, or W is a group M selected from a boron derivative or a metal function, and when q is 2 the other R^{1a} is R¹; with a compound Ar³-W¹, wherein W¹ is a halogen atom or a trifluoromethylsulfonyloxy group when W is a group M or W¹ is a group M when W is a halogen atom or a trifluoromethylsulfonyloxy group;


25 (d) to prepare a compound of formula (I) wherein R¹ is Ar³-Z and Z is O or S, reacting a compound of formula (V):



Formula (V)

30 wherein A, R², and q are as hereinbefore defined, one R^{1b} represents a group ZH and when q is 2 the other R^{1b} represents R¹; with a reagent serving to introduce the group Ar³;

(e) to prepare a compound of formula (I) where Y is a bond, reaction of a compound of formula (VI):



Formula (VI)

5 wherein R^1 , R^2 , q , Ar^1 and W are as hereinbefore defined, with a compound Ar^2-W^1 , wherein W^1 is a halogen atom or a trifluoromethylsulfonyloxy group when W is a group M , or W^1 is a group M when W is a halogen atom or a trifluoromethylsulfonyloxy group.

(f) interconversion of one compound of formula (I) to a different compound of formula (I) e.g. (i) alkylation of a compound (I) wherein R^2 represents hydrogen, (ii) conversion of one R^1 from alkoxy (e.g. methoxy) to hydroxy, or (iii) conversion of R^1 from hydroxy to sulfonyloxy, eg alkylsulfonyloxy or trifluoromethanesulfonyloxy; (iv) 10 conversion of a compound wherein Y represents S to a compound wherein Y is SO_2 or (v) conversion of Y from CO to CH_2 ;

(g) separation of *cis*- and *trans*- isomers of compounds of formula (I) by conventional methods; 15 and optionally thereafter forming a salt of formula (I).

6. A pharmaceutical composition comprising a compound of formula (I) as claimed in any of claims 1 to 4 or a physiologically acceptable salt thereof and a physiologically acceptable carrier therefor. 20

7. The use of a compound of formula (I) as claimed in any of claims 1 to 4 or a physiologically acceptable salt thereof in the manufacture of a medicament for the treatment of a condition which requires modulation of a dopamine receptor.

25 8. Use according to claim 7 wherein the dopamine receptor is a dopamine D_3 receptor.

9. Use according to claim 7 or claim 8 wherein a dopamine antagonist is required. 30

10. Use according to any of claims 7 to 9 wherein the condition is a psychotic condition.

11. Use according to claim 10 wherein the psychotic condition is schizophrenia. 35

12. A method of treating a condition which requires modulation of a dopamine

receptor which comprises administering to a subject in need thereof an effective amount of a compound of formula (I) as claimed in claim 1 or a physiologically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

Int. Application No.

PCT/EP 99/07763

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/12 C07D403/12 A61K31/55 C07D223/16 C07D471/04
 C07D413/12 C07D405/12 C07D409/12 C07D417/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	WO 98 50364 A (BRANCH CLIVE LESLIE ; JOHNSON CHRISTOPHER NORBERT (GB); STEMP GEOFF) 12 November 1998 (1998-11-12) the whole document	1,6-12
A	WO 98 06699 A (NASH DAVID JOHN ;STEMP GEOFFREY (GB); SMITHKLINE BEECHAM PLC (GB)) 19 February 1998 (1998-02-19) cited in the application the whole document	1,6-12
A	WO 97 43262 A (SMITHKLINE BEECHAM PLC ;STEMP GEOFFREY (GB); JOHNS AMANDA (GB)) 20 November 1997 (1997-11-20) cited in the application the whole document	1,6-12
	--- -/-- ---	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"Z" document member of the same patent family

Date of the actual completion of the international search

17 February 2000

Date of mailing of the international search report

23/02/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 eport,
 Fax: (+31-70) 340-3016

Authorized officer

Bosma, P

INTERNATIONAL SEARCH REPORT

Inter national Application No

PCT/EP 99/07763

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94 21628 A (MERCK SHARP & DOHME ;BROUGHTON HOWARD BARFF (GB); KULAGOWSKI JANUS) 29 September 1994 (1994-09-29) the whole document ——	1,6-12
A	EP 0 431 580 A (WARNER LAMBERT CO) 12 June 1991 (1991-06-12) cited in the application claims; examples ——	1,6-12
A	EP 0 007 070 A (SMITHKLINE CORP) 23 January 1980 (1980-01-23) ————	1,6-12

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 99/ 07763

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim 12
is directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compounds/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/07763

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9850364 A	12-11-1998	AU 7651898 A NO 995338 A	27-11-1998 02-11-1999
WO 9806699 A	19-02-1998	AU 4204697 A EP 0922035 A	06-03-1998 16-06-1999
WO 9743262 A	20-11-1997	AU 2897497 A BR 9709220 A CZ 9803632 A EP 0917530 A NO 985242 A PL 329803 A	05-12-1997 10-08-1999 16-06-1999 26-05-1999 09-12-1998 12-04-1999
WO 9421628 A	29-09-1994	AU 6215794 A US 5641787 A	11-10-1994 24-06-1997
EP 0431580 A	12-06-1991	US 4957921 A US 5047406 A AT 119522 T AU 628694 B AU 6770290 A CA 2031588 A DE 69017606 D DE 69017606 T DK 431580 T ES 2070253 T FI 905958 A IE 66740 B JP 2967943 B JP 3251568 A KR 171404 B NO 178262 B NZ 236335 A PT 96085 A, B ZA 9009787 A	18-09-1990 10-09-1991 15-03-1995 17-09-1992 13-06-1991 07-06-1991 13-04-1995 06-07-1995 24-07-1995 01-06-1995 07-06-1991 07-02-1996 25-10-1999 11-11-1991 01-02-1999 13-11-1995 26-05-1992 30-09-1991 26-08-1992
EP 0007070 A	23-01-1980	AR 222828 A AT 2267 T AT 19069 T AU 525789 B AU 4861379 A CA 1115271 A CS 213381 B DD 147355 A DK 278379 A, B, EP 0080012 A ES 482276 A ES 482281 A ES 482282 A FI 792125 A, B, GR 69675 A HU 179315 B IE 48655 B IE 48654 B IL 57532 A IT 1193190 B JP 1053271 B JP 55011584 A	30-06-1981 15-02-1983 15-04-1986 02-12-1982 07-02-1980 29-12-1981 09-04-1982 01-04-1981 08-01-1980 01-06-1983 01-04-1980 01-04-1980 01-04-1980 08-01-1980 07-07-1982 28-09-1982 03-04-1985 03-04-1985 31-08-1982 02-06-1988 13-11-1989 26-01-1980

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/07763

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0007070 A		MY 93387 A	31-12-1987
		NO 792267 A,B,	08-01-1980
		NZ 190745 A	23-01-1981
		PH 15754 A	18-03-1983
		PL 216908 A	11-08-1980
		PT 69855 A	01-08-1979
		SU 1029827 A	15-07-1983
		US 4265890 A	05-05-1981
		ZA 7902785 A	27-08-1980